



Russian Journal of Biological Research

Issued from 2014

E-ISSN 2413-7413
2023. 10(1). Issued one times a year

EDITORIAL BOARD

Bityukov Nikolai – Sochi State University, Russian Federation, Sochi, Russian Federation (Editor in Chief)

Davitashvili Magda – Telavi State University, Telavi, Georgia (Deputy Editor-in-Chief)

Khasanova Alfiya – Naberezhnye Chelny State Pedagogical University, Naberezhnye Chelny, Russian Federation

Marius Brazaitis – Lithuanian Sports University, Institute of Sports Science and Innovations, Kaunas, Lithuania

Volodin Vladimir – Komi Science Centre, Ural Branch of the Russian Academy of Sciences, Russian Federation

Journal is indexed by: **Cross Ref** (USA), **MIAR** (Spain), **Open Academic Journals Index** (USA).

All manuscripts are peer reviewed by experts in the respective field. Authors of the manuscripts bear responsibility for their content, credibility and reliability.

Editorial board doesn't expect the manuscripts' authors to always agree with its opinion.

Postal Address: 1717 N Street NW, Suite 1,
Washington, District of Columbia, USA
20036

Release date 25.12.23
Format 21 × 29,7/4.

Website: <https://rjbr.cherkasgu.press>
E-mail: office@cherkasgu.press

Headset Georgia.

Founder and Editor: Cherkas Global University
Order № B-15.

© Russian Journal of Biological Research, 2023

Russian Journal of Biological Research

2023

Is. 1

CONTENTS

Articles

Determining the Level of Soil Contamination Using an Oat Bioindicator A.O. Manaenkov, V.S. Bobrik	3
Evaluation of Antimalarial Activity of Methanolic Extract of <i>Carica Papaya (Carecaccaea)</i> Yellow Leaf in Mice M. Matawal Mankilik, Mhya, H. Daniel, P.I Wuyep	10
Ethnobotanical and Bioactive Characteristics of some <i>Sideritis</i> L. Taxa and Major Essential Oil Compounds Talip Şahin, Ö. Kılıç, T. Taşkın	17
Ethnobotanical Features and Pharmacological Studies of Some <i>Nepeta</i> L. Taxa T. Şahin, Ö. Kılıç	30

Copyright © 2023 by Cherkas Global University



Published in the USA
 Russian Journal of Biological Research
 Issued from 2014
 E-ISSN: 2413-7413
 2023. 10(1): 3-9

DOI: 10.13187/ejbr.2023.1.3
<https://rjbr.cherkasgu.press>



Articles

Determining the Level of Soil Contamination Using an Oat Bioindicator

Anton O. Manaenkov ^{a, *}, Vladimir S. Bobrik ^b

^a Academy of Management and Production, Moscow, Russian Federation

^b AHO OKB "Kristall", Moscow, Russian Federation

Abstract

The article presents the results of soil assessment using a bioindicator – germination of oats in room conditions. The task included sowing seeds and monitoring the dynamics of plant growth and development. The work of a team of two people ("B" and "A") was carried out on the territory of the Industry Agricultural Business Incubator, as well as at the home of each team member. As a result of the research, the following was noted. The shoots were unfriendly. Phytotoxicity coefficients showed that the cleanest point chosen for sampling was calculated to be the dirtiest of all. The seeds that grew best were those planted in the soil from the first (dirtiest) point; they had 100 % germination. As a result of seed germination, data was obtained that indicated that oats, with due attention, can grow equally in different soil conditions, perhaps this is due to the unpretentiousness of this plant or the insufficient distance between soil sampling points.

Keywords: oats, phytotoxicity, bioindicator, samples, above-ground part, underground part, soil.

1. Введение

Существует много способов оценки почвы. В данной работе она производится с помощью биоиндикатора, в роли которого выступил проращивание овса.

2. Обсуждение и результаты

Эксперимент

Опыт проходил у каждого из членов бригады на подоконнике. Первый опыт проводился в период с 01.11.19 по 12.11.19, но он оказался неудачным. Второй опыт начался 16 ноября в 23:00. Были посажены семена овса, на 4 сутки произошел практически дружный всход ростков, контейнеры с саженцами были помещены в разных точках: на освещенном участке, на письменном столе, а также в мало освещенном участке. Пробы почвы взяты из трёх точек ([Рисунок 1](#)).

* Corresponding author

E-mail addresses: manaenkov_ao@amp1996.ru (A.O. Manaenkov)



Рис. 1. Точки отбора проб



Рис. 2. Схема отбора проб почвы с точки № 2




Рис. 3. Схема отбора проб почвы с точек № 1 и № 3

1 точка – наиболее загрязненная точка, находящаяся рядом с дорогой, рядом проходит пешеходная дорожка, вымощенная плиткой.

2 точка – точка, обладающая умеренной антропогенной нагрузкой, выражающейся в удалении от проезжей части на 20 метров и относительной близостью лесного массива.

3 точка является наиболее чистым местом среди всех точек отбора почвы, так как она располагается рядом с Дендрологическим садом имени Р.И. Шредера ([Груздев, 2020](#)), ([Опекунова, 2016](#)).

Процесс отбора проб

На всех трёх точках сбор был одинаковым. На участке размером 50 × 50 см расчищалась земля от листьев, веток и камней до начала самого почвенного покрова. Далее подручными средствами (небольшими пластиковыми стаканами и лопатками) производилось выкапывание земли до глубины приблизительно 15 см. Собранный грунт помещался в пакет, чтобы донести до места обработки.

Далее в течение недели происходила просушка почвы (на бумаге формата А3) так, чтобы лишняя влага испарилась, и земля оказалась сухой, после этого почва очищалась от лишних примесей и перемешивалась до равномерной консистенции и далее производилась посадка.

Первый опыт проращивания был неудачный. Овес не пророс, причина – обильное количество воды в почве.

Второй опыт: после просушки почва снова просеивалась, убрались лишние семена и проростки, оставшиеся после прошлого опыта, далее посев происходил повторно.

Взяв по 6 (на каждую точку по 2) контейнеров-стаканов на 2 человека («В» и «А») из бригады, производилось рассаживание семян. Предварительно в контейнеры положили по одной салфетке, чтобы задерживать почву, которая вымывалась бы при поливе растений. Далее, просеяв почву, началось выкладывание ее слоем в 4-5 см. Далее в каждый контейнер клалось по 10 семян (в сумме 120), которые немного присыпались тонким слоем сухой земли (0,3–0,5 мм). Затем следовала не обильный полив, чтобы семена прижились. Засев происходил 16 ноября в 22:00. В течение следующих двух недель отслеживалось состояние, периодически поливая или же брызгая с пульверизатора контейнеры, и уже 20 ноября произошел всход первых ростков (Скупченко, Соколова, 2009), (Гелашвили, 1995), (Чеснокова, Чугай, 2008).



Рис. 4. Проростки второго посева



Рис. 5. Проба на пятый день опыта

По итогам произведенных испытаний были получены данные, которые сведены в Таблицах 1–5. В Таблицах 1–2 вписаны значения измерений в формате: длина ростка от семечка/длина корешка. Все измерения даны на последний день опыта. На основе Таблиц 1–5 произведены выводы и данные по которым сведены в Таблицах 6–7.

Таблица 1. Пробы проращивания овса (опыт «В»)

Точка	Данные (надземная часть / подземная часть, в см)									
1.1	11,5 / 3,4	11 / 10	13,4 / 2,2	15 / 3,8	20 / 4	13,6 / 6,5	15,2 / 5,2			
1.2	20,7 / 3,4	18,5 / 1,9	15,7 / 3,8	13,5 / 2,9	14,4 / 1,8	22,9 / 3,6	18,2 / 3,8	18,7 / 4,5	13,1 / 2,3	
2.1	16,7 / 4,6	5,6 / 1,2	12,2 / 2,4							
2.2	18,2 / 4,8	19,5 / 3,2	17,9 / 3,5	16,8 / 4,7	19,7 / 5,2	21,2 / 1,6	23,6 / 3,9			
3.1	18,7 / 3,4	19,5 / 2,5	21,3 / 2,9	21,7 / 4,1	8,6 / 1,2	13,8 / 4,6	16,4 / 2,7	18,9 / 3,1		
3.2	18,9 / 3,9	22,2 / 2,9	20,3 / 3,1	17,5 / 2,6	20,8 / 3,6					

Таблица 2. Пробы проращивания овса (опыт «А»)

Точка	Данные (надземная часть / подземная часть, в см)									
1.1	24 / 3,5	22 / 2,8	8,3 / 1,4	18 / 1,5	24,2 / 5,4	23 / 4,2	7,7 / 2,8			
1.2	13 / 2,8	22 / 4,5	20 / 5,6	21,8 / 6,5	24 / 2,8	23 / 4,7	6 / 2,9	22,5 / 2,8		
2.1	15,5 / 2	18,4 / 2,8	14,3 / 2,1	18 / 3,9	13,2 / 1,7	15 / 3,8				
2.2	23,5 / 2	21,5 / 3	20,7 / 7	23,1 / 4,2	21,3 / 3,5	19,5 / 3,6	16 / 2,1	17 / 2,9	22 / 1,5	22,5 / 3,3
3.1	25 / 3,5	21 / 4,2	19,6 / 3,2	16,7 / 2,9	20,7 / 3,7	20,3 / 5,2				
3.2	11,5 / 3,5	11,8 / 3,1	22,1 / 3,6	22,5 / 2,8	17,9 / 2,6					

Таблица 3. Средние значения вершков и корешков по точкам, в сантиметрах

Точка	«В»		«А»	
	Надземная часть	Подземная часть	Надземная часть	Подземная часть
1.1	12,94	3,15	18,17	3,08
1.2	17,3	3,1	18,04	2,9
2.1	11,5	2,73	15,73	2,72
2.2	19,0	4,3	20,71	3,31
3.1	19,56	3,06	20,55	3,78
3.2	19,94	3,22	17,16	3,12

Таблица 4. Всхожесть по дням

День	20.11	21.11	22.11	23.11	24.11	25.11	26.11
«В»							
1.1	0	2	4	5	5	6	7
1.2	1	3	3	4	6	8	9

День	20.11	21.11	22.11	23.11	24.11	25.11	26.11
2.1	0	0	0	1	2	3	3
2.2	0	2	3	5	6	6	7
3.1	1	2	3	5	6	6	8
3.2	0	0	2	3	4	5	5
«А»							
1.1	0	3	4	5	5	7	7
1.2	1	1	4	6	7	7	8
2.1	0	1	3	4	5	5	6
2.2	2	4	6	8	9	10	10
3.1	2	3	4	4	6	6	6
3.2	2	3	4	4	4	5	5

Таблица 5. Сравнительная таблица

Всхожесть проростков (%)	Характеристика проростков	Оценка загрязнения
Менее 20	Мелкие и уродливые	Сильное
20-60	Тонкие и короткие	Среднее
60-90	Почти нормальной длины	Слабое
90-100	Ровные и крепкие	Отсутствует

Таблица 6. Всхожесть ростков в процентах

	Точки					
	1.1	1.2	2.1	2.2	3.1	3.2
«В»	70	90	30	70	80	50
«А»	70	80	60	100	60	50

Таблица 7. Оценка загрязнения почвы

	Точки					
	1.1	1.2	2.1	2.2	3.1	3.2
«В»	Слабое / Отсутствует		Среднее / Слабое		Слабое / Среднее	
«А»	Слабое		Слабое / Отсутствует		Слабое / Среднее	

Проанализировав Таблицы 7, можно сделать вывод о том, что первая точка является наиболее чистой, вторая – среднезагрязненной, а третья самой загрязненной из всех.

$$\Phi_3 = \frac{L_K - L}{L_K} * 100, \% \quad (1)$$

где Φ_3 – коэффициент фитотоксичности; L_K – длина вершка или корешка на контроле (чистая точка), L – длина вершка/корешка на загрязненной или среднезагрязненной точке.

$$\begin{aligned} \Phi_3(2\text{т}''\text{В}''\text{надз.}) &= \frac{15,25-15,12}{15,25} * 100 = 0,85\% \\ \Phi_3(2\text{т}''\text{В}''\text{подз.}) &= \frac{3,515-3,125}{3,515} * 100 = 11,09\% \\ \Phi_3(3\text{т}''\text{В}''\text{надз.}) &= \frac{19,75-15,12}{19,75} * 100 = 23,44\% \\ \Phi_3(3\text{т}''\text{В}''\text{подз.}) &= \frac{3,14-3,125}{3,14} * 100 = 0,48\% \\ \Phi_3(2\text{т}''\text{А}''\text{надз.}) &= \frac{18,22-18,105}{18,22} * 100 = 0,63\% \\ \Phi_3(2\text{т}''\text{А}''\text{подз.}) &= \frac{3,015-2,99}{3,015} * 100 = 0,83\% \\ \Phi_3(3\text{т}''\text{А}''\text{надз.}) &= \frac{18,855-18,105}{18,855} * 100 = 3,98\% \\ \Phi_3(3\text{т}''\text{А}''\text{подз.}) &= \frac{3,45-2,99}{3,45} * 100 = 13,33\% \end{aligned}$$

Таблица 8. Сводная таблица по фитотоксичности, %

	«В»	«А»
2 точка – надземная часть	0,85	0,63
2 точка – подземная часть	11,09	0,83
3 точка – надземная часть	23,44	3,98
3 точка – подземная часть	0,48	13,3

Исходя из результатов расчетов коэффициентов фитотоксичности, можно сделать вывод, что коэффициент на второй точке меньше, чем на третьей. Это свидетельствует о том, что третья точка отбора проб более загрязненная (Белюченко и др., 2014; Манаенков и др., 2020).

3. Заключение

В результате проведенных исследований было отмечено следующее.

Всходы были недружные. Коэффициенты фитотоксичности показали, что самая чистая точка, выбранная для отбора проб, является по результатам расчетов самой грязной из всех. Лучше всего росли те семена, посаженные в почву с первой (самой грязной) точки, дали 100 % всхожесть.

Точки отбора проб были расположены примерно в одной местности – на территории отраслевого бизнес-инкубатора. В итоге проращивания семян были получены данные, которые свидетельствовали о том, что овёс при должном внимании может произрастать одинаково в разных почвенных условиях, возможно, это связано с неприхотливостью данного растения или недостаточной удаленностью между точками отбора проб почвы.

Литература

Белюченко и др., 2014 – Белюченко И.С., Федоненко Е.В., Смагин А.В. Биомониторинг состояния окружающей среды: Учебн. пособие / Под ред. ред. И.С. Белюченко, Е.В. Федоненко, А.В. Смагин. Краснодар: КубГАУ, 2014. 153 с.

Гелашвили, 1995 – Гелашвили Д.Б. Экологический мониторинг. Ч. 1. Методы биомониторинга: Учебн. пособие. Нижн. Новгород: Изд-во ННГУ, 1995. 190 с.

Груздев, 2020 – Груздев В.С. Биоиндикация состояния окружающей среды. М.: ИНФРА-М, 2020. 160 с.

Манаенков и др., 2020 – Манаенков А.О., Шаховская К.Д., Науменко Н.О. Определение экологического состояния озера Бездонного в парке «Серебряный Бор» Методом Вудивисса / Географическое изучение территориальных систем: Сб. мат. XIV Всеросс. научн.-практ. конф. студентов, аспирантов и молодых учёных. Пермь, 2020. С. 184-18.

Опекунова, 2016 – Опекунова М.Г. Биоиндикация загрязнений: Учебн. пособие. СПб.: Изд-во СПбГУ, 2016. 300 с.

Скупченко, Соколова, 2009 – Скупченко В.Б., Соколова Л.О. Биоиндикация окружающей среды: Учебн. пособие. СПб.: Изд-во СПбГЛТУ, 2009. 72 с.

Чеснокова, Чугай, 2008 – Чеснокова С.М., Чугай Н.В. Биологические методы оценки качества объектов окружающей среды: Учебн. пособие. В 2-х ч. Ч. 2. Методы биотестирования. Владимир: Изд-во ВлГУ, 2008. 92 с.

References

Belyuchenko i dr., 2014 – Belyuchenko, I.S., Fedonenko, E.V., Smagin, A.V. (2014). Biomonitoring sostoyaniya okruzhayushchei sredy [Biomonitoring of the environment: Textbook. allowance]: Uchebn. posobie. Pod. red. I.S. Belyuchenko. Krasnodar: KubGAU, 153 p. [in Russian]

Chesnokova, Chugai, 2008 – Chesnokova, S.M., Chugai, N.V. (2008). Biologicheskie metody otsenki kachestva ob"ektov okruzhayushchei sredy [Biological methods for assessing the quality of environmental objects]: Uchebn. posobie. V 2-kh ch. Ch. 2. Metody biotestirovaniya. Vladimir: Izd-vo VIGU, 92 p. [in Russian]

Gelashvili, 1995 – Gelashvili, D.B. (1995). *Ekologicheskii monitoring. Ch. 1. Metody biomonitoringa* [Environmental monitoring. Part 1. Methods of biomonitoring]: Uchebn. posobie. Nizhn. Novgorod: Izd-vo NNGU, 190 p. [in Russian]

Gruzdev, 2020 – Gruzdev, V.S. (2020). *Bioindikatsiya sostoyaniya okruzhayushchei sredy* [Bioindication of the state of the environment]. M.: INFRA-M, 160 p. [in Russian]

Manaenkov i dr., 2020 – Manaenkov, A.O., Shakhovskaya, K.D., Naumenko, N.O. (2020). *Opreделение ekologicheskogo sostoyaniya ozera Bezdonnogo v parke «Serebryani Bor» Metodom Vudivissa* [Determination of the ecological state of Lake Bezdonoynoye in the Serebryany Bor park using the Woodiwiss method]. *Geograficheskoe izuchenie territorial'nykh sistem: Sb. mat. XIV Vseross. nauchn.-prakt. konf. studentov, aspirantov i molodykh uchenykh*. Perm'. Pp. 184-18. [in Russian]

Opekunova, 2016 – Opekunova, M.G. (2016). *Bioindikatsiya zagryaznenii* [Bioindication of pollution]: Uchebn. posobie. SPb.: Izd-vo SPbGU, 300 p. [in Russian]

Skupchenko, Sokolova, 2009 – Skupchenko, V.B., Sokolova, L.O. (2009). *Bioindikatsiya okruzhayushchei sredy* [Bioindication of the environment]: Uchebn. posobie. SPb.: Izd-vo SPbGLTU, 72 p. [in Russian]

Определение уровня загрязнения почвы с помощью биоиндикатора овса

Антон Олегович Манаенков ^{a,*}, Владимир Сергеевич Бобрик ^b

^a Академия управления и производства, Москва, Российская Федерация

^b АХО ОКБ «Кристалл», Москва, Российская Федерация

Аннотация. В статье представлены результаты оценки почвы с помощью биоиндикатора – проращиванием овса в комнатных условиях. В задачу входило проведение посева семян, отслеживание динамики роста и развития растений. Работа бригады из двух человек («В» и «А») проводилась на территории Отраслевого аграрного бизнес-инкубатора, а также в домашних условиях каждого члена команды. В результате проведенных исследований было отмечено следующее. Всходы были недружные. Коэффициенты фитотоксичности показали, что самая чистая точка, выбранная для отбора проб, является по результатам расчетов самой грязной из всех. Лучше всего росли те семена, посаженные в почву с первой (самой грязной) точки, дали 100 % всхожесть. В итоге проращивания семян были получены данные, которые свидетельствовали о том, что овёс при должном внимании может произрастать одинаково в разных почвенных условиях, возможно, это связано с неприхотливостью данного растения или недостаточной удаленностью между точками отбора проб почвы.

Ключевые слова: овёс, фитотоксичность, биоиндикатор, пробы, надземная часть, подземная часть, почва.

* Корреспондирующий автор

Адреса электронной почты: manaenkov_ao@amp1996.ru (А.О. Манаенков)

Copyright © 2023 by Cherkas Global University



Published in the USA
 Russian Journal of Biological Research
 Issued from 2014
 E-ISSN: 2413-7413
 2023, 10(1): 10-16

DOI: 10.13187/ejbr.2023.1.10
<https://rjbr.cherkasgu.press>



Evaluation of Antimalarial Activity of Methanolic Extract of *Carica Papaya* (*Carecaccaea*) Yellow Leaf in Mice

Mary Matawal Mankilik^{a,*}, Mhya H. Daniel^b, Ponjul Wuyep^a

^a Department of Biochemistry, Faculty of Basic Medical Sciences, University of Jos, Nigeria

^b Department of Biochemistry, Faculty of Basic Medical Sciences, Abubakar Tafawa Balewa University Bauchi, Nigeria

Abstract

Malaria still remains a public health problem in developing countries. Malaria is one of the most prevalent diseases as people still rely on traditional medicine as source of treatment for this disease. Malaria causes mortality and morbidity with social economic impact in developing countries where the burden is high. Reports showed that the high global health challenges is partly due to multidrug resistance *P. falciparum* develop on existing and available antimalarial drugs and that has spur the search of alternate treatment with low or less side effect. The purpose of this study is to evaluate the antimalarial potential of methanolic extract of *Carica papaya* yellow leaf on animal model. In vivo screening for antimalarial drug discovery is of the recommended stream line process for new compound in path from drug discovery to developing. The Rane's curative method of established infection was employed in vivo for assessing antimalarial activity. Swiss albino mice of both sexes weighing between 23-27 g and aged 6 weeks were infected with 1×10^7 *P. berghei* (NK-65) RBC/ml intraperitoneally and were treated with various dose (100, 200 and 400mg/kg b.wt) of *C. papaya* yellow leaf methanolic extract. Acute oral toxicity test was employed using OECD method. The mice treated with 400 mg/kg b.wt of *C. papaya* yellow leaf extract showed significant ($P < 0.05$) antimalarial activity. In acute oral toxicity result showed that the maximum tolerated dose was found to be 5000 mg/kg body weight. *C. papaya* leaf extract showed antimalarial activity and study has validated its use by locals in the treatment of malaria in most developing countries. Recommendations are made for the isolation and identification of bioactive substance for possible drug development.

Keywords: Antimalarial activity, *P. berghei* (NK-65), infected mice, *C. papaya* yellow leaf.

1. Introduction

Malaria is a serious pathogenic disorder, which causes mortality and morbidity with social and economic impact most especially in developing countries where the burden of malaria is very high. According to WHO 2021, there were an estimated 241 million malaria cases and 627,000 malaria related cases worldwide which represent 14 million more case in 2021 compound to 2019, and 69,000 more death. Which were linked to disruption to the provision of malaria prevention, diagnosis and treatment drug the pandemic. This disease has positioned about 3.3 billion people at risk. The recommended preventive drugs include a combination of sulfadoxine, pyrimethamine and amodiaquine, while the therapeutic strategy includes use of first line drugs – artemisinin

* Corresponding author

E-mail addresses: Mankilikma@unijos.edu.ng (M. Mary Mankilik)

compound therapy in areas where *P. falciparum* is endemic and chloroquine (CQ), in the areas where it is still efficient.

The high global challenges developed by existing and partly available antimalarial drug due to multidrug resistance *P. falciparum* has led to the urgent need in search for newer treatments to eradicate malaria most especially in developing countries.

Although the available synthesized drugs may have side effects, indigenous drugs are better alternate sources for the treatment of malaria due to low or less side effect. The resistance in malaria parasite is believed to have emerged through mutation in the active sites of drug targets or from biochemical changes in drug receptors.

The plant *C. papaya* L (English Paw-Paw; Hausa: Gwanda; Marghi; Ron: Mban) is a large tree-like plant belonging to the *Caricaceae* family, it is commonly called "Paw-Paw". *C. papaya* is widely grown in tropical and subtropical lowlands region. Among the developing countries, India, Indonesia, Mexico, Nigeria and Thailand are the largest papaya grown countries (Wimalawansa, 1981).

It contains pawpaw, chymopapain vegetable pepsin (Wimalawansa, 1981). The presence of papaya endopeptidase II and *Carica papaya* endopeptidase IV were also reported. Protein, carpaine were also found to be present (Gills, 1992), leaves contain Carposide and seed contain Myrosinase carian and Sinigrin glycoside (Otskia et al., 2010). Phytochemicals present includes tanins, flavonoid, saponin, anthroquinones, steroids reducing sugars, cardinolides and phenolics (Owoyele et al., 2008).

2. Material and methods

Collection and authentication of plant materials

The yellow leaf of *C. papaya* were collected from university permanent site Naraguta Campus in April 2021 and was authenticated by a taxonomist at the Department of Plant Science and Biotechnology, Faculty of Natural Science University of Jos, Nigeria and specimen with voucher number JUHN21000418 was deposited in the departmental herbarium.

Preparation of the yellow leaf extract

After the collection of the yellow leaf water to remove any dirt and foreign particles and then shade dried. The dried yellow leaves were then crushed and grounded to get a powder form and weighed the powdered material (50 g) was weighed and extracted with 300 ml of methanolic for 72 hrs as described by Mankilik et al., 2021a, Mankilik et al., 2021b the solution obtained was sieved and extract kept in a desiccator to evaporate. It was then kept in a clean sterile bottle for further use.

Chemicals

All chemicals and reagents used were of analytical grade procured from various certified local and international suppliers.

Acute oral toxicity studies

A safe oral dose of the extract was determined by accurate oral toxicity test of Organization of Economic Cooperation and Development (OECD, 2008) as per 425 guidelines as described by Mankilik et al., 2021a. The animals were observed for any sign of toxicity OECD, 2008.

Experimental animals

Swiss albino mice weighing 18-30 g of either sex were used in the study. All the animals were housed for a week in a ventilated standard environment condition. The animals were used in this study were obtained from animal house unit of Pharmacology Department, Faculty of Pharmaceutical Sciences, University of Jos. The experiment was approved by the ethical committee Animal experimental unit, Department of Pharmacology and experiment were conducted as per guidelines of Institutional Animal Care and Use Committee (IACU) in collaboration with Office of Laboratory Animal Welfare office of laboratory animal (OLAW) with approval number F17-00379.

Maintenance of *P. berghei*

The chloroquine (CQ) sensitive rodent malarial parasite *P. berghei* (NK-65) strain was purchased from the National Institute of Medical Research, Ibadan, Oyo State, Nigeria. The blood stage parasite was maintained in an adult donor mice by serial blood passage and mice with 20-30 % rising parasitaemia was anaesthetized using chloroform dapped in a cotton wool. The blood

collected in heparinized tubes contain 0.5 % trisodium citrate buffer and adjusted with 0.9 % physiological saline so that each 0.2 ml aliquate contain 1×10^{-7} infected red blood cell/ml of blood.

Antimalarial test – Rane's curative assay

In the curative assay on Day 0 (D_0) mice were intraperitoneally infected with 1×10^{-7} *P. berghei* infected erythrocytes followed by random division of into four groups of 3 mice, each group 72 hrs after observation, the groups 1-3 orally received the plant extract of 200, 300 and 400 mg/kg bwt/day respectively. The group 4 (negative control) and group 5 (positive control) received 0.2 ml of vehicle and CQ phosphate 25 ml/kg b.wt/day respectively. The animals were dosed accordingly once daily for 5 consecutive days (D_2 - D_7). On D_8 , the blood parasiteamia in the all the groups were determined by using Giemsa stain. The mean survival of the treatment group were. Arithmetically determined by calculating the average survival time starting from day of infection to 30 days (D_0 - D_{29}), in the Rane's test, the body weight was measured 3 hrs between infection on day 0 and consecutively from D_3 – D_8 during treatment period to establish the effect of plant drug on malarial mice (Ryley, Peters, 1970).

Determination of mean survival time

Mortality in each group was monitored daily from the time of parasite inoculation to death in the treatment and control groups throughout the study period. In the curative assay (Rane's test) the survival was observed for 30 days (D_0 – D_{29}). The Mean Survival Time (MST) of each group was calculated as:

$$\text{MST} = \frac{\text{Sum of survival time of all mice on a group}}{\text{total number of mice in a group}}$$

Statistical data analysis

Each experiment was done in triplicate and data's were expressed as mean $\pm$ standard error on mean. Statistical significance was determined by one way analysis variance (ANOVA). The $P \leq 0.05$ from percentage parasiteamia were considered statistically significant when compared to control group.

3. Results

Phytochemical screening

The phytochemical screening of *C. papaya* yellow leaf methanolic extract reveals the presence of flavonoid, tannins, balsam, phenols and resins (Table 1).

Acute toxicity

The result of the toxicity study was depicted in Table 2. The extract was found to be safer up to 5000mg/kg b.wt.

Body weight

Result of Figure 1 showed that body weight of mice in the control group (CQ = 25 mg/kg) significantly increased, compared to negative control and other treatment groups. Dose dependent decrease was seen in treatment group (400 > 300 > 200).

PCV

Extracts did not ameliorate reduction in PCV level (Figure 2) reduction in PCV treatment group is dose is dependent as standard drug treatment group prevented reduction in PCV.

Antiplasmodial activity of plant extract and chloroquine

The result showed that the percentage increase in mice treated with extract at different doses on day 2 (D_2) while the parasiteamia of the mice treated with CQ 10 mg/kg did not increase compound compared to to those treated with *C. papaya* extract on day 2. In the chloroquine treated group, there was significant reduction in *P. berghei* to non-detectable level (D_8 – D_{10}). (Figure 3) there was significant ($P < 0.05$) decrease in parasiteamia level of mice infected with *P. berghei* and treated with 400mg/kg of *C. papaya* yellow leaf extract on Day 10 (D_{10}).

Mean Survival Time

All extract prolonged survival time, with 400 mg/kg b.wt of extract surviving being standard drug experimental period compared to the infected untreated (Table 3).

Table 1. Phytochemical profile of the Yellow Leaves Methanolic Extract On *Carica papaya*

PHYTOCHEMICAL	RESULT	
Alkaloids	+	
Flavonoids	+	
Tannins	+	
Saponins	-	
Terpenes and steroids		+
Cardiac glycosides	-	
Balsams	+	
Carbohydrates		-
Phenols	+	
Resins	+	

KEY

- = absent


+ = present

Table 2. Acute toxicity of plant extract in mice with behavioral changes

Plant	Behavioral changes
<i>C. papaya</i> Yellow leaf methanolic extract	2000 mg/kg b.wt 5000 mg/kg b.wt No behavioral changes or mortality, except hair erection, no other symptoms were observed, No signs of toxicity and mortality was observed.

Table 3. Mode of Survival Time (MST) of Swiss albino mice

GROUPS	SURVIVAL RATE
200 mg/kg b. wt	18 days
300 mg/kg b. wt	16 days
400 mg/kgb.wt	Lived beyond experimental period
Infected/Untreated	13 days
25 mg/kgb. wt of chloroquine	Lived beyond experimental period

**Fig. 1.** Effect of yellow leaves methanolic extract of *Carica papaya* on body weight of *Plasmodium berghei* mice




Fig. 2. Effect of yellow leaves methanolic extract of *Carica papaya* on PCV of *Plasmodium berghei* mice




Fig 3. In vivo antiplasmodial activity of yellow leaves extract of *Carica papaya* against *Plasmodium berghei* infected mice

4. Discussion

Malaria remain the most vital pathogen disease. The rodent malaria parasite *P. berghei* is reported to be analogous to human malaria parasites in most essential aspect of structures, physiology life-cycle and model for the investigation of biology of malaria parasites, parasite – host interaction, vaccine development and drug testing (Fidock et al., 2004).

In toxicity test of yellow leaf methanol extract of *C. papaya* showed no visible toxic effects up to 5000 mg/kg b.wt. This study is similar to our previous study on the aqueous and methanolic extract of *A. aspera* shoot (Mankilik et al., 2021a; Mankilik et al., 2021b).

In the antiplasmodial study, the chloroquine sensitive *P. berghei* (NK-65) strain was used to check the efficacy of *C. papaya* yellow leaf methanolic extract along with conventional antimalarial

agent chloroquine. Hence in the antimalarial screening the drug is expected to prevent body weight loss under parasiteamia. The crude extract of *C. papaya* at various doses prevented weight loss induced with increased in parasiteamia level. In our study there was steady decrease in weight by extract treated group, except for dose 400 mg/kg b.wt where steady increase was recorded. Decrease in weight has been associated with decreased food intake, disturbed metabolic function and likely hypoglycaemia (Atkinson et al., 2000; Bashar et al., 2012).

The plant extract was effective in curative assay exhibiting above 50 % antiplasmodial activity. In our studies there was a remarkable increase of parasiteamia level by infected untreated groups of mice with remarkable decrease in chloroquine treated group from day 8 to a non-detectable level. The extract treated groups also reduced *P. berghei* levels compared with infected untreated. The present study was also comparable to our previous antiplasmodial study reported (Mankilik et al., 2021a; Mankilik et al., 2021b) on *A. aspera* L.; *Vernoma amgadalina* (Longdet et al., 2021); *Eucalyptus cameldulensis* ethylacetate extract (Longdet et al., 2020) with array of positive approaches for medicinal plants in malaria. The methanolic extract revealed the presence of bioactive component contained in the plant such as, terpenoids, alkaloids, flavonoids, tannins and phenol. Proving that the may be the major constituent for antimalarial activity. This activity could be attributed to the presence of good concentration of active compounds in higher doses (Zaruwa et al., 2018).

The observation of mean survival time in days revealed that the drug treated group prolonged the survival time of mice compared to the untreated group. The chloroquine group animals' survival beyond the experimental period (D₂₉) compared to negative control which displayed 13 days of survival. At higher doses of 400 mg/kg b.wt of *C. papaya*, survival days were beyond the experimental period, while at dose 200 mg/kg and 300 mg/kg b.wt prolonged survival of mice for 18 and 16 days respectively. These could be due to the reduced parasiteamia could and overall reduction in pathologic effect of the parasite strain on the study mice and thus agreed with the report that bioactive compounds of medicinal plants that prolonged survival time greater than twelve days in considered as active (Ural et al., 2014). This study is consistent with previous study (Mankilik et al., 2021a; Mankilik et al., 2021b).

From our study, there was remarkable decline in PCV levels of mice in all treatment group from D₂-D₁₀ compared to the standard control. The low PCV observed in the infected and untreated groups may be as a result of heamolysis due to growing infection. This is in tandem with report that induced reduction in PCV levels during malaria in rodent occurred approximately 48 hrs post infection (Mace et al., 2015). The destruction of RBC either by parasite multiplication or spleen reticuloendothelial cell action in the pressure of many abnormal RBC stimulate the spleen to produce many phagocytes which could lead to aneamia *P. berghei* infected mice (Chinchilla et al., 1998). *Carica papaya* displayed antimalarial effect this has ability to reduce the erythrocytes stages (merozoites, schizonts and gametocytes) of *P. berghei* and hence justify its traditional usage of malaria remedy. This is in tandem with previous research that *C. papaya* L. extracted from leaf was reported having antidiengue, anti-inflammatory and antimalarial (Charan et al., 2016; Pandey et al., 2016; Ansari, 2016).

5. Conclusion

The study showed that yellow leaf methanolic extract of *Carica papaya* leaf could possess useful phytochemicals and may be a potential candidate for isolation of antimalarial compound and may provide scientific support for its use in traditional medicine. Extract is safe as no sign of mortality or morbidity was seen.

6. Conflict of interest

The authors declare that they have no competing interest.

7. Acknowledgements

The authors would like to acknowledge the Department of Biochemistry and staff of Animal House unit.

Ethics and consent approval to participate:

The study was approved by the institute of Animal Ethical Committee, University of Jos, Ethical clearance obtained with approval number UJ/FPS/F17-00379.

References

- Akhilar, Vijayalakshmi, 2015 – Akhilar, S., Vijayalakshmi, N.G. (2015). Phytochemical studies on *Carica papaya* leaf juice. *IJPSR*. 6(2): 880-883.
- Ansari, 2016 – Ansari, R.M. (2016). Extract of *Caricap papaya* L. leaves: standardizing its use in dengue fever. *Indian Journal of Pharmacology*. 48: 338-339.
- Charan et al., 2016 – Charan, J., Saxena, X., Goyal, J.P., Yasobant, S. (2016). Efficacy and safety of *Carica papaya* leaf extract in the dengue: A systemic review and meta-analysis. *International Journal of Applied & Basic Medical Research*. 6:249-254.
- Evans, Evans, 2002 – Evans, W.C., Evans, T. (2002). Pharmacognosy (1st Edition) N.B Saunders Company LTD, London 191-393.
- Fidock et al., 2004 – Fidock, D.A., Rosenthal, P.J., Croft, S.L., Brun, R., Nwaka, S. (2004). Antimalarial drug discovery; efficacy models for compound screening. *Nature Reviews Drug Discovery*. 3(6): 509-520. DOI: 10.1038/nrd1416
- Gills, 1992 – Gills, L.S. (1992). Edition Ethnomedicinal uses of plants in Nigeria. University of Benin Press, Edo State. 263.
- Longdet, 2019 – Longdet, I.Y., Mankilik, M.M., Idoko, E.D. (2019). Anti-plasmodial activity of *Eucalyptus camaldulensis* in mice. *African Journal of Biomedical Research*. 22: 303-307.
- Longdet, 2019 – Longdet, I.Y., Mankilik, M.M., Offiku, O.M., Idoko, E.D. (2019). Evaluation of the antiplasmodial activities of diethyl ether leaf extract of experimental mice. *Russian Journal of Biological Research*. 6(1): 8-15.
- Mankilik et al., 2021a – Mankilik, M.M., Longdet, I.Y., Luka, C.D. (2021). Antiplasmodial Activity of Methanolic Extract of *Achyranthes aspera* Shoot against *Plasmodium berghei* Infection in Albino Mice. *Russian Journal of Biological Research*. 8(1): 27-35. DOI: 10.13187/ejbr.2021.1.27
- Mankilik et al., 2021b – Mankilik, M.M., Longdet, I.Y., Luka, C.D. (2021). Evaluation of *Achyranthes aspera* shoot extract as an alternative therapy for malaria. *Journal of Basic and Applied Zoology*. 82(14). DOI: <https://doi.org/10.1186/s4136-021-00211-4>
- Mutabingwa, 2003 – Mutabingwa, T.K. (2003). Treating, malaria pregnancy in Africa. Rostawhate Doctor Africa. 26066-73.
- Olliaro, 2003 – Olliaro, P.L., Tailor, W.R.J. (2003). Antimalarial compounds from bench to bedside. *Journal of Experimental Biology*. 206: 3753-3759.
- Otsukia et al., 2010 – Otsukia, N., Dangh, N.H., Kamagaik, E., Kondoc, A., Iwataa, S., Morimoto, C. (2010). Aqueous extract of *Carica papaya* extracts anti-tumor activity and immunomodulatory effects. *J Ethnopharm*. 127: 760-767.
- Owoyele, 2008 – Owoyele, B.V., Adebukola, O.M., Funmilayo, A.A., Soladoye, A. (2008). Anti-inflammatory activities of ethamine extract of *Carica papaya* leaves. *Inflammopharm*. 16: 168-175.
- Pandey et al., 2016 – Pandey, S., Cabot, P.J., Shaw, P.N., Hewavitharana, A.K. (2016). Anti-inflammatory and immunomodulatory properties of *C. papaya*. *Journal of Immunotoxicology*. 13: 590-602.
- Peters, 1982 – Peters, W. (1982). Antimalarial drug resistance: An increasing problem. *Brazilian Medical Build*. 32: 187-192.
- Ryley, 1995 – Ryley, J., Peters, W. (1995). The antimalarial activity of some quinole esters. *Ann Trop Medcinal Parasitology*. 84(22): 209-222.
- Schuster, 2001 – Schuster, B.G. (2001). Demonstrating the validity of natural products as anti-infective drugs. *Journal of Alternative and Complementary Medicine*. 7(1): 573-582.
- Sirima, 2003 – Sirima, S.B., Sawadogo, R., Moran, A.C., Konate, A., Diarra, A. et al. (2003) failure of chloroquine chewoprophylaxis program to adequately prevent malararia during pregnancy in Kompela distribution. *Burkinafaso Clinical Infectious Diseases*. 36: 1374-1386.
- Wimalawansa, 1981 – Wimalawansa, S.J. (1981). Papaya – the treatment of chronic infected ulcers. *Medical Journal*. 26: 129-132.

Copyright © 2023 by Cherkas Global University



Published in the USA
 Russian Journal of Biological Research
 Issued from 2014
 E-ISSN: 2413-7413
 2023. 10(1): 17-29

DOI: 10.13187/ejbr.2023.1.17
<https://rjbr.cherkasgu.press>



Ethnobotanical and Bioactive Characteristics of some *Sideritis* L. Taxa and Major Essential Oil Compounds

Talip Şahin ^{a, *}, Ömer Kılıç ^b, Turgut Taşkın ^c

^a Department of Biology, Institute of Science, Adıyaman University, Turkey

^b Department of Pharmaceutical Botanic, Faculty of Pharmacy, Adıyaman University, Turkey

^c Department of Pharmacognosy, Faculty of Pharmacy, Marmara University, Turkey

Abstract

Turkey with a huge and rich variety of flora reserves lots of medical, aromatic and bioactive properties plants within its structure. In Turkey public medical science applications are encountered extensively and Turkey is one of the leading countries in the trade of medicinal and aromatic plants because of its geographical location, difference climate conditions, plant diversity and agricultural potential. As in whole world and in Turkey, use of plants existing in the natural flora for different purposes in traditional medicine; as spice, tea, insecticide, resin, veterinary cure, glue, essential oil, beverage, beekeeping, landscape and cosmetic industry. On the other hand, this important ethnobotanical knowledges are facing a danger of being lost by contemporary urbanisation and other negative factors.

The local population has long utilized the taxa of the genus *Sideritis* L., as herbal tea and in traditional medicine. Most of *Sideritis* taxa have significant biological activity, the biological activities and traditional usage of the *Sideritis* taxa are far better understood in recent years but still needs attention. The majority of *Sideritis* taxa exhibit a variety of biological functions and are highly interesting for future research on their pharmacological characteristics and potential for pharmacy and medicine.

In this study, it is aimed to review major essential oil compound, usage of traditional knowledge and biological activities of some *Sideritis* taxa in Turkey.

Keywords: *Sideritis*, Ethnobotany, Medicinal plant, Aromatic plant, Traditional use, Biological activity, Essential oil.

1. Introduction

Sideritis L. genus is in the Lamiaceae family. Lamiaceae is a family include economically important grasses, shrubs and rarely in tree form. *Sideritis*, *Salvia*, *Thymus*, *Mentha*, *Origanum*, *Phlomis*, *Nepeta*, *Teucrium*, *Rosmarinus*, *Lavandula*, *Thymbra*, *Satureja*, *Ballota*, *Stachys*, *Ajuga*, *Prunella*, *Melissa*, *Lamium*, *Marrubium* are most known genres of Lamiaceae; which have many aromatic and medicinal importance taxa. While Lamiaceae members naturally grow in many parts of the world, especially in the gene center of Mediterranean region, Labiatae is represented by very few species in cold regions.

* Corresponding author

E-mail addresses: talipsahin34@gmail.com (T. Şahin), omerkilic77@gmail.com (Ö. Kılıç), ttaskin237@gmail.com (T. Taşkın)

The Lamiaceae family is one of the rich and important cosmopolitan families of Angiosperms, represented by 224 genera and approximately 5600 species in the world. According to the Flora of Turkey, there are approximately 45 genera and more than 850 taxa of which more than 235 are endemic. In addition, Lamiaceae members are in the first place among the plants traded in Turkey (Kılıç, 2014).

Members of Lamiaceae have a unique aromatic odor and are aromatic plants, and these fragrances they carry are due to essential oils. Due to have abundant essential oils, they have been used in medicine, cuisine, landscaping, pharmacy and perfumery for a long time. In addition, it is known that aromatic compounds, which are abundant in Lamiaceae members, affect the mood and health of individuals intensely and are used extensively in phytotherapy and aromatherapy. The Lamiaceae family is the family of plants most known and studied for their essential oils all over the world (Lee et al., 2011).

The genus *Sideritis* is one of the genus with the highest endemism rate in Turkey, and approximately 1/3 of the *Sideritis* taxa are endemic for Turkey. Due to its high endemism rate (79.5 %), Turkey is one of the gene centers of this genus. In Turkey, the species belonging to the genus *Sideritis* are known by different names among the people (Kılıç, 2002).

In this review study, major essential oil compound, usage of traditional knowledge and biological activities of some *Sideritis* taxa in Turkey are compiled.

2. Results and discussion

Usage properties of some *sideritis* taxa

Medicinal aromatic plants are generally used as a source of tea, spice, folk medicine and source of essential oil, and *Sideritis* members are ethnobotanically important and economically valuable plants used for these purposes. The first information about the use of *Sideritis* taxa as medicinal plants dates back to the first century. In many country *Sideritis* taxa are used as herbal tea. In addition, *Sideritis* taxa are usage as herbal tea, many taxa are used in the traditional medicine for treatment of different diseases and health problems (Başer, 2000).

Sideritis taxa are widely used as folk medicine and herbal tea in Turkey ethnobotany. Commonly used because of its nervous system stimulant, anti-inflammatory, antispasmodic, carminative, analgesic, sedative, antitussive, stomachic and anticonvulsant effects. Members of the *Sideritis* genus are also widely used in the treatment of colds and gastrointestinal diseases (Table 1).

Sideritis taxa are also widely used in the aromatherapy and in phytotherapy, because they contain high rates of fragrant oils and commonly used aromatic substances. For this reason, some members of the *Sideritis* genus, which have high economic value. *Sideritis* species are often used in the treatment of inflammation and gastrointestinal disorders. These activities are based on the antioxidant properties of terpenoids and flavonoids in its composition (Koleva et al., 2002).

Sideritis albiflora, *Sideritis congesta*, *Sideritis libanotica*, *Sideritis perfoliata* and *Sideritis sipylea* have also been reported to be used as analgesics in stomach pain (Özaydın et al., 2005). *S. vulcanica* is consumed as an antipyretic and as an antipyretic among the people, and it is also ground and inhaled as a powder against headache (Arslan, 1999). *Sideritis libanotica* as a sedative, in colds, in the treatment of skin wounds, as an astringent. It has been reported that *S. montana* is used as a cough suppressant (Altundağ, Öztürk, 2011).

According to another study, *Sideritis perfoliata* is used in folk medicine for colds, bronchitis, stomach disorders. It has been noted that *Sideritis trojana* is also used in stomach disorders, abdominal pain, laxative, kidney diseases and sore throat (Bulut, Tuzlacı, 2015). *Sideritis trojana* was used as a folk remedy for digestive problems and colds (Polat, Satıl, 2012).

In another study, *Sideritis leptoclada* was used as a sedative in the treatment of colds and shortness of breath. It has been noted that *Sideritis montana* is used as a folk remedy in the treatment of stomach ailments (Kılıç et al., 2021). It has been reported that *Sideritis albiflora* is used as a diuretic, diabetic, relieving stomach and respiratory ailments (Sarı et al., 2010). *S. scardica* for calming down after work and *Sideritis condensata* is used traditionally for stress relief (Bruno et al., 2002).

Many *Sideritis* taxa are used for treatment of flu and colds, respiratory diseases, diseases of the digestive system, diseases of the urinary system, diseases of the cardiovascular system and anemia. In addition pharmacological activity of *Sideritis* members can be listed as; antimicrobial

and antiviral activity, antioxidant activity, anti-inflammatory activity, gastroprotective activity, pharmacologic activities on the central nervous system diseases, cytotoxic activity and other activities (Aneva et al., 2019).

Table 1. Used part, ethnobotanical uses, bioactivity and major essential compound of some *Sideritis* taxa.

Botanical name	Location of the Plant	Used Part	Preparation method	Ethnobotanical uses	Reported bioactivities	Major essential oil compound
<i>S. athea</i>	Bingöl, Kazdağı/Balıkesir, Edremit/Balıkesir	Aerial parts, flowers	Infusion	Common cold, flu, stomach ailments (Polat et al., 2011; Kalankan et al., 2015) (Polat, Satıl, 2012; Şahin, Kılıç, 2022)	Antimicrobial, Antidepressant, Ability to memorize (Kılıç et al., 2003; Öztürk et al., 1996)	Carvacrol (Küçük et al., 2021)
<i>S. bilgeriana</i>	Aladaglar/Nigde	Aerial part	Infusion	Heartburn, cold, hemorrhoids (Özdemir, Alpınar, 2015; Şahin, Kılıç, 2022)	Antimicrobial, Antioxidant (Dulger et al., 2006; Iscan et al., 2005; Aydın et al., 1996)	β-pinene (Iscan et al., 2005)
<i>S. brevidens</i>	Eğirdir/Isparta, Erdemli/Mersin, Gölhisar/Burdur, Fethiye/Muğla	Unspecified	Unspecified	Common cold (Başer et al., 2006)	Antioxidant, Antimicrobial, Antifungal (Tunalier et al., 2004; Dülger et al., 2005)	β-pinene (Kirimer et al., 1999)
<i>S. caesarea</i>	Pinarbasi/Kayseri	Aerial parts	Fresh herb is ingested. boiled in water, after filtering drunk as tea	Stomach ailments, ulcer (Gürbüz et al., 2005; Şahin, Kılıç, 2022)	Antifungal, Antimicrobial, Antioxidant, Antiulcerogenic, Antioxidative (Gürbüz et al., 2005; Dulger et al., 2006; Sağdıç et al., 2008; Aşkun et al., 2008; Çelik, Kaya, 2011)	β-caryophyllene (Günbatan et al., 2017)
<i>S. condensata</i>	The Middle Aegean Region	Aerial parts	Decoction	Food (Kargioğlu et al., 2010)	Antimicrobial, Antioxidant (Dülger et al., 2005; Özkan et al., 2001; Özkan et al., 2005; Güvenç et al., 2010; Özcan et al., 2005)	β-caryophyllene (Kirimer et al., 1996)
<i>S. erythrantha</i> var. <i>cedretorum</i>	Bozyazı/Mersin	Aerial parts	Infusion, gargle, mixture, spice	Cold, flu, pharyngitis, blurred vision, pleasure, as tea (Sargin, 2015)	Antimicrobial, Antioxidant (Köse et al., 2010)	α-pinene (Tabanca, 2001)
<i>S. erythrantha</i> var. <i>erythrantha</i>	Bozyazı/Mersin, Sütçüler/Isparta	Aerial parts, leaf, flowers	Infusion, gargle, mixture (Pickles, jam i.e.), spice	Cold, flu, pharyngitis, blurred vision, pleasure, as tea, antiseptic, germicidal, nausea, cold, asthma, desiccant, night sweats, spinal cord discomfort, blood purifier, gas reliever, breast milk booster (Sargin, 2015; Demirel, 2021)	Antimicrobial, Antioxidant, Antiviral (Köse et al., 2010; Altundağ et al., 2011; Altundağ, Aslım, 2011; Özkan et al., 2005; Yesilada, 2023)	α-pinene (Başer, Kirimer, 2018)
<i>S. germanicopolitana</i>	Kisecik/Karaman	Flowers	Infusion	For tea purpose (Yücel et al., 2016)	No studies found	Myrcene (Kirimer et al., 1992)
<i>S. lanata</i>	Bozyazı/Mersin	Aerial parts	Infusion	Kidney renewing, pleasure, as tea (Sargin, 2015)	Antidepressant, Antimicrobial (Uğur et al., 2005)	Spathulenol (Kirimer, 2000)
<i>S. leptoclada</i>	Acıpayam/Denizli	Aerial parts	Infusion	Sedative (Bulut et al., 2017)	Antimicrobial, Antioxidant (Saraç, Uğur, 2007; Askun et al., 2009; Güvenç et al., 2005)	Germacrene D (Deveci, 2019)

<i>S. libanotica</i> Subsp. <i>microchlamys</i>	Gölbasi/Adiyaman, Tut/Adiyaman, Birecik/Şanlıurfa, Adiyaman (centre), Yaslica/Şanlıurfa	Aerial part	Infusion	Stomach relaxing, fatigue relieving, pleasant, cold, flu, shortness of breath, cough (Güldaş, 2009; Furkan, 2016; Balos, Hakan, 2007; Abak, 2018)	Antioxidant, Antimicrobial, Cytotoxicity, Anticholinesterase (Atas et al., 2019)	β -caryophyllene (Kirimer et al., 2004)
<i>S. libanotica</i> subsp. <i>linearis</i>	Kürecik/Malatya, Midyat/Mardin, Andırın/Kahramanmaraş	Aerial parts, leaf	Infusion, decoction	Food, cold, for tea purpose (Kargioğlu et al., 2010; Akgül et al., 2018; Demirci, Özhatay, 2012; Tevent, 2020)	Antioxidant, Insektisit, Anti-inflammatory, Antimicrobial, Antiproliferative (Ayhan, 2008; Demirtaş et al., 2009; Yeşilada, Ezer, 1989; Ezer et al., 1994; Ezer et al., 1995)	α -bisabolol (Erbaş, Fakir, 2012)
<i>S. libanotica</i> subsp. <i>libanotica</i>	Sarıveliler/Karman, Hatay (centre), Yeşildere/Karman	Flowers, aerial part	Infusion, decoction	For tea purpose, appetizer, carminative, sedative (Bağcı et al., 2016; Güzel et al., 2015; Akdag, Dogu, 2016)	Antioxidant (Güvenc et al., 2005)	Sideridiol (Formisano et al., 2015)
<i>S. libanotica</i> subsp. <i>violascens</i>	Sarıveliler/Karman, Yeşildere/Karman	Flowers	Infusion	As tea (Bağcı et al., 2016; Akdag, Dogu, 2016)	Antimicrobial (Yılmaz, 2013)	β -caryophyllene (Kirimer et al., 2004)
<i>S. libanotica</i> subsp. <i>kurdica</i>	East Anatolia, Artuklu/Mardin, Bozova/Şanlıurfa, Şanlıurfa (centre), Gölpınar/Şanlıurfa	Leaf, flowers, aerial parts	Infusion, decoction, diabetes, a pinch is consumed as raw.	Sedative, colds, skin wounds, astringent, for tea purpose, gargle, diabetes (Abak, 2018; Altundag, Ozturk, 2011; Akan, Ayaz, 2015; Fidan, 2018; Kılıç, 2019; Oymak, 2018; Şahin-Fidan, 2018)	Antidepressant (Öztürk, 1996)	α -pinene (Kirimer et al., 2004)
<i>S. montana</i> subsp. <i>montana</i>	Kırklareli (centre)	Aerial parts	Decoction	Cold, flu, cough (Kültür, 2011)	Anticancer, Antioxidant (Venditti et al., 2016)	Germacrene D (Kirimer et al., 2000)
<i>S. perfoliata</i>	Kazdağı/Bahkesi r, Bayramiç/Çanak kale, Acıpayam/Deniz li, Andırın/Kahraman maraş, Sütçüler/Isparta, Bigadiç/Bahkesi r, Bergama/Izmir	Aerial parts, leaf, flowers	Infusion	As tea, cold, bronchitis, stomach ailments, diuretic (Saçlı, Akalin, 2001)	Antimicrobial, Antidepressant, Anticancer, Ability to memorize Antioxidant, Anti- inflammatory, Reducing blood pressure, Hypoglycaemic, Antioxidant, Antimicrobial, Cytotoxic, Anticholinesterase (Ezer et al., 1994; Kirimer et al., 2008; Atalay, 2014; Çarıkçı, 2020)	Limonene (Kirimer et al., 2008)
<i>S. phrygia</i>	Yahyalı- Pınarbaşı/Kayseri, Develi/Kayseri	Flowers	Infusion	As tea (Çelikel, 2002)	Antioxidant (Öztürk et al., 1996; Güvenc et al., 2005)	Limonene (Özer-Sağır, 2016)
<i>S. pisidica</i>	Yörenler- Elmalı/Antalya, Dumanlı-	Leaf	Infusion, externally as porridge	Tonic, as a tea, for stomachaches (Yeşilada et al.,	Antimicrobial, Anti- inflammatory, Antioxidant,	Caryophyllene (Özer-Sağır, 2016)

	Beyşehir/Konya			1993) (Yeşilada et al., 1995)	Anticholinesterase, Antitumor (Dülger et al., 2005; Yeşilada, Ezer et al., 1994; Özer-Sağır, 2016; Deveci et al., 2017)	
<i>S. rubriflora</i>	Bozyazı/Mersin	Aerial parts	Infusion, gargle	Cold, flu, pharyngitis, pleasure, as tea (Demirel, 2021; Bulut et al., 2017; Demirci, Özhatay, 2012; Bulut, Tuzlacı, 2015; Kocabaş, Gedik, 2016; Tanaydın, 2021; Arasan, 2022)	Antimicrobial, Antifeedant, Antifungal, (Dulger et al., 2006; Bondi et al., 2000)	β -pinene (Kirimer et al., 1999)
<i>S. scardica</i> subsp. <i>scardica</i>	Kırklareli (centre),	Aerial parts	Decoction	Bronchitis, cough, cold, flu (Sargin, 2015; Şahin, Kılıç, 2022)	Antioxidant (Tunalier et al., 2004)	b-pinene (Solomou et al., 2019)
<i>S. sipylea</i>	Sarıgöl/Manisa	Aerial parts	Infusion	Foot odor, haircare, pleasure, cold, flu, as tea (Kültür, 2007)	Antimicrobial, Antioxidant (Dulger et al., 2006; Güvenc et al., 2005; Loğoğlu et al., 2006; Aligiannis et al., 2001; Gergis et al., 1991)	β -myrcene (Maškovic et al., 2023)
<i>S. stricta</i>	Aladaglar/Nigde	Aerial parts	Infusion	Cold, flu (Akgül et al., 2016; Sarı et al., 2010)	Antimicrobial, Alzheimer's disease, Parkinson's disease, Anti-inflammatory, Analgesic (Küpeli et al., 2007; Kılıç, 2006; Turkmenoglu et al., 2015; Küpeli et al., 2007)	β -pinene (Sattar, 1995)
<i>S. syriaca</i> subsp. <i>nusairiensis</i>	Kahramanmaraş (centre)	Aerial parts	Unspecified	Cold (Güzel et al., 2015; Kocabaş, Gedik, 2016; Altay et al., 2015; Varlıbaş-Odunkıran, 2020; Özer, Türkmen, 2019)	Antiviral, Antimicrobial, Antioxidant, Analgesic, Antiinflammatory (Sattar, 1995; Koutsaviti et al., 2013; Goulas et al., 2014; Karapandzova et al., 2013; Menghini et al., 2005)	α -pinene (Kirimer et al., 2004)
<i>S. trojana</i>	Kazdağı/Bahkesir, Bayramiç/Çanakale, Edremit/Balıkesir	Herbs, aerial parts	Infusion	As tea, stomach ailments, Abdominal pain, laxative, kidney ailments, sore throat, cold (Şahin, Kılıç, 2022; Bulut, Tuzlacı, 2009)	Cytotoxic, Antimicrobial, Anti-helicobacter, Insecticidal, Antioxidant, Antifungal, (Kılıç et al., 2003; Dulger et al., 2006; Kirimer et al., 2008; Kılıç et al., 2003; Kirmızibekmez et al., 2017; Aslan et al., 2006; Kirmızibekmez et al., 2012)	β -pinene (Kirimer et al., 2008)
<i>S. vulcanica</i>	Arıcak/Elazığ	Aerial parts	Infusion	Appetizer, germicidal, anti-inflammatory, carminative, urinary tract infections, Cold (Polat, Satıl, 2012; Saçlı, Akalın, 2001; Bulut, Tuzlacı, 2015)	Antioxidant (Tunalier et al., 2004)	α -pinene (Başer, Kirimer, 2018)

3. Conclusion

In Turkey ethnobotany the domestic people used *Sideritis* taxa for preparing beverages like herbal tea, and also for the treatment of various diseases and health problems, as wound healing and for some blood disorders.

Most information available is about the use of *Sideritis* taxa in the case of respiratory diseases, especially cold and flu. Many *Sideritis* taxa also have high antibacterial and antiviral activities. These activities are supposed to be due mostly to the terpenes, iridoids, some flavonoids, sterols and essential oils. The traditional use of many species of *Sideritis* genus for treatment of digestive system diseases could be linked to the inflammatory activity, due to the flavonoids.

Studies of the biological activities of *Sideritis* taxa (Figure 1) are dominated by these focused on the antioxidant and antibacterial activity of the plant extracts. Still, the predominant studies are of screening type, with emphasis on the selecting of appropriate species, extraction method and testing material, while the studies on the mechanism of action of the active substances are scarce.




Fig. 1. Biological activities of some *Sideritis* taxa in human body (Aneva et al., 2019).

In conclusion, *Sideritis* taxa offers a wide source in different research fields and these taxa are provides a wide range of possibilities that can be found in further scientific studies like; phytochemistry, phytoteraphy, pharmacology, pharmacognosy, aromateraphy, toxicology etc.

4. Conflict of interest

The authors declare that there are no conflicts of interest.

References

- Abak, 2018** – Abak, F. (2018). Şanlıurfa İli Lamiaceae (Ballıbabagiller) Familyasının Florası Bazı Taksonların Fitokimyasal Ve Etnobotanik Özellikleri. Recep Tayyip Erdoğan Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Doktora Tezi.
- Akan, Ayaz, 2015** – Akan, H., Ayaz, H. (2015). Gölpınar (Şanlıurfa-Türkiye) mesire yeri florası ve etrafındaki köylerin etnobotanik özellikleri. *Bağbahçe Bilim Dergisi*. 2(3):19-56.
- Akdag, Dogu, 2015** – Akdag, T., Dogu, S. (2015). The Medical Plants Of Karaman-Yesildere Village And It's Surroundings. *International Journal of Agriculture and Environmental Research*. 2(5): 1214-1225.
- Akgul et al., 2018** – Akgul, A., Akgul, A., Senol, S.G., Yildirim, H., Secmen, O., Dogan, Y. (2018). An ethnobotanical study in Midyat (Turkey), a city on the silk road where cultures meet. *Journal of Ethnobiology and Ethnomedicine*. 14:12. DOI: 10.1186/s13002-017-0201-8
- Akgül et al., 2016** – Akgül, G., Yılmaz, N., Celep, A., Celep, F., Çakılcıoğlu, U. (2016). Ethnobotanical purposes of plants sold by herbalists and folk bazaars in the center of Cappadocia (Nevşehir, Turkey). *Indian Journal of Traditional Knowledge*. 15(1): 103-108.
- Aligiannis et al., 2001** – Aligiannis, N., Kalpoutzakis, I., Chinou, B., Mitakou, S. (2001). Composition and antimicrobial activity of the essential oils of five taxa of *Sideritis* from Greece. *J Agric Food Chem*. 49: 811-5.
- Altay et al., 2015** – Altay, V., Karahan, F., Sarcan, Y.B., İlçim, A. (2015). An ethnobotanical research on wild plants sold in Kırkhan district (Hatay/Turkey) herbalists and local markets. *Biological Diversity and Conservation*. 8(2): 81-91
- Altundağ et al., 2011** – Altundağ, S., Aslım, B., Öztürk, S. (2011). In vitro antimicrobial activities of essential oils from *Origanum minutiflorum* and *Sideritis erythraea* subs. *erythraea* on phytopathogenic bacteria. *J Essent Oil Res*. 23: 4-8.
- Altundağ, Aslım, 2011** – Altundağ, S., Aslım, B. (2011). Effect of some endemic plants essential oils on bacterial spot of tomato. *J Plant Pat*. 93(1): 37-41.
- Altundağ, Öztürk, 2011** – Altundağ, E., Öztürk, M. (2011). Ethnomedicinal studies on the plant resources of east Anatolia, Turkey. *Procedia Soc. Behav. Sci*. 19:756-777.
- Aneva et al., 2019** – Aneva, I., Zhelev, P., Kozuharova, E., Danova, K., Nabavi, S.F., Behzad, S. (2019). Genus *Sideritis*, section *Empedoclia* in southeastern Europe and Turkey – studies in ethnopharmacology and recent progress of biological activities. *DARU Journal of Pharmaceutical Sciences*. 27: 407-421.
- Arasan, 2022** – Arasan, Ş. (2022). Kozak Yaylası (Bergama) Ve Çevresinde Etnobotanik Araştırmalar. Ege Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Doktora Tezi.
- Arslan, 1999** – Arslan, K. (1999). *Sideritis vulcanica* Hub.-Mor. Üzerine Anatomik, Morfolojik ve Korolojik Çalışmalar. Balıkesir Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Eğitimi Anabilim Dalı. Yüksek Lisans Tezi.
- Askun et al., 2009** – Askun, T., Tumen, G., Satıl, F., Ates, M. (2009). Characterization of the phenolic composition and antimicrobial activities of Turkish medicinal plants. *Pharm Biol*. 47: 563-71.
- Aslan et al., 2006** – Aslan, I., Kılıç, T., Gören, A., Topçu, G. (2006). Toxicity of acetone extract of *Sideritis trojana* and 7-epicandiciol, 7-epicandiciol diacetate and 18-acetylsideroxol against stored pests *Acanthoscelides obtectus* (Say), *Sitophilus granarius* (L.) and *Ephesia kuehniella* (Zell). *Ind Crops Prod*. 23: 171-6.
- Aşkun et al., 2008** – Aşkun, T., Tumen, G., Satıl, F., Kılıç, T. (2008). Effects of some Lamiaceae species methanol extracts on potential mycotoxin producer fungi. *Pharm Biol*. 46 (10-11): 688-694.
- Atalay, 2014** – Atalay, B. (2014). Kazdağları'nda Yetişen Lamiaceae Familyasının Bazı Türlerinin Biyolojik Aktiviteleri. Balıkesir Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.
- Atas et al., 2019** – Atas, M., Eruygur, N., Sozmen, F., Ergul, M., Ergul, M., Akpulat, H.A., Ucar, E. (2019). Evaluation of Various Biological Activities of Endemic *Sideritis libanotica* Extracts. *Not Sci Biol*. 11(2): 210-217.
- Ayaz, 2008** – Ayaz, A. (2008). *Sideritis hololeuca* Boiss. & Heldr. apud Bentham ve *Sideritis libanotica* Labill. subsp. *violascens* ekstrelerinin antibakteriyel aktivitelerinin belirlenmesi. Selçuk Üniversitesi, Yüksek Lisans Tezi.

- Aydın et al., 1996 – Aydın, S., Ozturk, Y., Beis, R., Baser, K.H.C. (1996). Investigation of *Origanum onites*, *Sideritis congesta* and *Satureja cuneifolia* essential oils for analgesic activity. *Phytother Res.* 10: 342-4.
- Ayhan, 2008 – Ayhan, B. (2008). *Sideritis Libanotica Linearis* Bitkisinin Sekonder Metabolitlerinin Saflastırılması, Karakterizasyonu Ve Bazı Biyolojik Aktivitelerinin İncelenmesi. Gaziosmanpaşa Üniversitesi, Fen Bilimleri Enstitüsü, Kimya Anabilim Dalı, Yüksek Lisans Tezi.
- Bağcı et al., 2016 – Bağcı, Y., Erdoğan, R., Doğu, S. (2016). Sarıveliler (Karaman) ve Çevresinde Yetişen Bitkilerin Etnobotanik Özellikleri. *Nisan.* 42 (1): 84-107.
- Balos, Akan, 2007 – Balos, M.M., Akan, H. (2007). Zeytinbahçe - Akargay (Birecik, Şanlıurfa) Arasında Kalan Bölgenin Etnobotanik Özellikleri. *S. Ü. Fen Ed. Fak. Fen Derg.* 29: 155-171.
- Başer et al., 2006 – Başer, K.H.C., Tümen, G., Malyer, H., Kırmır, N. (2006). Plants used for common cold in Turkey. *Proceedings of ICEB* (2005). Pp. 133-137.
- Başer, 2000 – Başer, K.H.C. (2000). Uçucu yağların parlak geleceği. Tıbbi ve Aromatik Bitkiler Bülteni: 15, Anadolu Üniversitesi Tıbbi ve Aromatik Bitki ve İlaç Araştırma Merkezi, Eskişehir.
- Başer, Kırmır, 2018 – Başer, K.H.C., Kırmır, N. (2018). Essential oils of Anatolian Lamiaceae - An update. *Nat. Volatiles & Essent. Oils.* 5(4): 1-28.
- Bondi et al., 2000 – Bondi, M., Bruno, M., Piozzi, F., Baser, K.H.C., Simmonds, M. (2000). Diversity and antifeedant activity of diterpenes from Turkish species of *Sideritis*. *Biochem Syst Ecol.* 28: 299-303.
- Bruno et al., 2002 – Bruno, M., Rosselli, S., Pibiri, I., Kilgore, N., Lee, K.H. (2002). Anti-HIV agents from the ent-kaurane diterpenoid linearol. *J Nat Prod.* 65: 1594-7.
- Bulut et al., 2017 – Bulut, G., Haznedaroğlu, M.Z., Doğan, A., Koyu, H., Tuzlacı, E. (2017). An Ethnobotanical Study Of Medicinal Plants In Acıpayam (Denizli-Turkey). *Herbal Medicine.* DOI: <http://dx.doi.org/10.1016/j.hermed.2017.08.001>
- Bulut, Tuzlacı, 2009 – Bulut, G., Tuzlacı, E. (2009). Folk Medicinal Plants Of Bayramiç. *J. Fac. Pharm.* İstanbul. 40.
- Bulut, Tuzlacı, 2015 – Bulut, G., Tuzlacı, E. (2015). An Ethnobotanical Study of Medicinal Plants in Bayramiç (Çanakkale-Turkey). *Marmara Pharmaceutical Journal.* 19: 268-282.
- Çarıkçı, 2020 – Çarıkçı, S. (2020). Antioxidant and Anticholinesterase Properties of *Sideritis perfoliata* subsp. *athoa* (Papan. & Kokkini) Baden and *Sideritis trojana* Bornm. Teas from Mount Ida-Turkey and Their Phenolic Characterization by LCMS/MS. *JOTCSA.* 7(2): 617-634.
- Charami et al., 2008 – Charami, M.T., Lazari, D., Karioti, A., Skaltsa, H., Hadjipavlou-Litina, D., Souleles, C. (2008). Antioxidant and antiinflammatory activities of *Sideritis perfoliata* subsp. *perfoliata* (Lamiaceae). *Phytother Res.* 22: 450-4.
- Çarıkçı et al., 2007 – Çarıkçı, S., Çöl, Ç., Kılıç, T., Azizoğlu, A. (2007). Diterpenoids from *Sideritis tmolea* P.H. Davis. *Rec Nat Prod.* 1(4): 44-50.
- Çelik, Kaya, 2011 – Çelik, I., Kaya, M.S. (2011). The antioxidant role of *Sideritis caesarea* infusion against TCA toxicity in rats. *Br J Nutr.* 105: 663-8.
- Çelikel, 2002 – Çelikel, Ö. (2002). Kayseri ve çevresinde halk tarafından kullanılan bitkilerin yöresel adları ve kullanım amaçları. Erciyes Üniversitesi, Yüksek Lisans.
- Demirci, Özhatay, 2012 – Demirci, S., Özhatay, N. (2012). An Ethnobotanical Study In Kahramanmaraş (Turkey); Wild Plants Used For Medicinal Purpose In Andırın, Kahramanmaraş. *Turk J. Pharm. Sci.* 9(1): 75-92.
- Demirel, 2021 – Demirel, O. (2021). Sütçüler (Isparta)' De Etnobotanik Bir Çalışma. Isparta Uygulamalı Bilimler Üniversitesi, Lisansüstü Eğitim Enstitüsü, Orman Mühendisliği Anabilim Dalı, Yüksek Lisans Tezi.
- Demirtaş et al., 2009 – Demirtaş, I., Şahin, A., Ayhan, B., Tekin, S., Telci, I. (2009). Antiproliferative effects of the methanolic extracts of *Sideritis libanotica* Labill. subs. *linearis*. *Rec Nat Prod.* 3(2): 104-109.
- Deveci et al., 2017 – Deveci, E., Tel-Çayan, G., Yıldırım, H., Duru, M.E. (2017). Chemical composition, antioxidant, anticholinesterase and anti-urease activities of *Sideritis pisdica* Boiss. & Heldr. endemic to Turkey. *Marmara Pharmaceutical Journal.* 21(4): 898-905.
- Deveci et al., 2019 – Deveci, E., Tel-Çayan, G., Usluer, Ö., Duru, M.E. (2019). Chemical Composition, Antioxidant, Anticholinesterase and Anti- Tyrosinase Activities of Essential Oils of Two *Sideritis* Species from Turkey. *Iranian Journal of Pharmaceutical Research.* 18(2): 903-913.

Doğanoğlu et al., 2006 – Doğanoğlu, Ö., Gezer, A., Yücedağ, C. (2006). Göller Bölgesi-Yenişarbademli Yöresi'nin Önemli Bazı Tıbbi Ve Aromatik Bitki Taksonları Üzerine Araştırmalar. *Süleyman Demirel Üniversitesi Fen Bilimleri Enstitüsü Dergisi*. 10: 66-73.

Dülger et al., 2005 – Dülger, B., Ugurlu, E., Aki, C., Suerdem, T.B., Camdeviren, A., Tazeler, G. (2005). Evaluation of antimicrobial activity of some endemic Verbascum, Sideritis, and Stachys species from Turkey. *Pharm Biol*. 43: 270-4.

Erbaş, Fakir, 2012 – Erbaş, S., Fakir, H. (2012). Türkiye'nin Batı Akdeniz Yöresinde doğal olarak yetişen dağ çayı (Sideritis libanotica Labill. subsp. linearis (Benth) Bornm) ve bayır kekiği (Origanum sipyleum L.) türlerinin uçucu yağ oranları ve bileşenlerinin belirlenmesi. *SDU Faculty of Forestry Journal*. 13: 119-122.

Ezer et al., 1994 – Ezer, N., Usluer, G., Güneş, İ., Erol, K. (1994). Antibacterial activity of some Sideritis species. *Fitoterapia*. LXV(6): 549-550.

Ezer et al., 1995 – Ezer, N., Akçoş, Y., Rodriguez, B., Abbasoğlu, U. (1995). Sideritis libanotica subsp. linearis (Benth) Bornm., den elde edilen iridoit heteroziti ve antimikrobiyal etkisi. *Hacettepe Üniversitesi Eczacılık Fakültesi Dergisi*. 15(1):15-21.

Formisano et al., 2015 – Formisano, C., Oliviero, F., Rigano, D., Arnold N.A., Senatore, F. (2015). Comparative Chemical Composition and Antioxidant Properties of the Essential Oils of three Sideritis libanotica Subspecies. *Natural Product Communications*. 10 (6): 1075-1078.

Furkan, 2016 – Furkan, M.K. (2016). Adıyaman İlinde Yetişen Bazı Bitkilerin Etnobotanik Özellikleri. Adıyaman Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.

Gergis et al., 1991 – Gergis, V., Spiliotis, V., Argiriadou, N., Poulos, C. (1991). Relation between the antimicrobial activity and the chemical composition of the essential oil of Sideritis sipylea Boiss. *Fragrance J*. 6: 93-5.

Goulas et al., 2014 – Goulas, V., Exarchou, V., Kanetis, L., Gerothanassisa, I. (2014). Evaluation of the phytochemical content, antioxidant activity and antimicrobial properties of mountain tea (Sideritis syriaca) decoction. *J Funct Foods*. 6: 248-58.

Gülbatan et al., 2017 – Günbatan, T., Demirci, B., Gürbüz, İ., Demirci, F., Gençler-Özkan, A.M. (2017). Comparison of Volatiles of Sideritis caesarea Specimens Collected from Different Localities in Turkey. *Natural Product Communications*. 12(10): 1639-1642.

Gültaş, 2009 – Gültaş, H. (2009). Adıyaman İlinde Etnobotanik Değeri Olan Bazı Bitkilerin Kullanım Alanlarının Tespiti. Fırat Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.

Gürbüz et al., 2005 – Gürbüz, I., Özkan, A.M., Yesilada, E., Kutsal, O. (2005). Anti-ulcerogenic activity of some plants used in folk medicine of Pinarbasi (Kayseri, Turkey). *Journal of Ethnopharmacology*. 101: 313-318.

Güvenc et al., 2005 – Güvenc, A., Houghton, P.J., Duman, H., Coskun, M., Sahin, P. (2005). Antioxidant activity studies on selected Sideritis species native to Turkey. *Pharm Biol*. 43: 173-7.

Güvenç et al., 2010 – Güvenç, A., Okada, Y., Akkol, E., Duman, H., Okuyama, T., Çalı, I. (2010). Investigations of anti-inflammatory, antinociceptive, antioxidant and aldose reductase inhibitory activities of phenolic compounds from Sideritis brevibracteata. *Food Chem*. 118: 686-92.

Güzel et al., 2015 – Güzel, Y., Güzelsemme, M., Miski, M. (2015). Ethnobotany Of Medicinal Plants Used In Antakya: A Multicultural District In Hatay Province Of Turkey. *Journal of Ethnopharmacology*. DOI: <http://dx.doi.org/10.1016/j.jep.2015.07.042>

Iscan et al., 2005 – Iscan, G., Kirimer, N., Kurkcuoğlu, M., Baser, K.H.C. (2005). Composition and antimicrobial activity of the essential oils of two endemic species from Turkey: Sideritis cilicica and Sideritis bilgerana. *ChemNat Compd*. 41: 679-82.

Kalankan et al., 2015 – Kalankan, G., Özkan, Z.C., Akbulut, S. (2015). Medicinal and Aromatic Wild Plants and Traditional Usage of Them in Mount Ida (Balıkesir/Turkey). *Journal of Applied Biological Sciences*. 9(3): 25-33.

Kan et al., 2018 – Kan, Y., Kan, A., Ayran, İ., Çelik, S.A. (2018). Essential Oil Yield and compositions of endemic mountain tea (Sideritis libanotica Labill. ssp. linearis (Benth) Bornm. end Sideritis bilgerana P.H. Davis) cultivated in Konya ecological conditions of Turkey. *Int J Agric Environ Food Sci*. 2(1): 204-205.

Kargioğlu et al., 2010 – Kargioğlu, M., Cenkeci, S., Serteser, A., Konuk, M., Vural, G. (2010). Traditional Uses Of Wild Plants In The Middle Aegean Region Of Turkey. *Hum Ecol*. 38: 429-450.

- Kılıç et al., 2003 – Kılıç, T., Yıldız, Y.K., Gören, A.C., Tümen, G., Topçu, G. (2003). Phytochemical analysis of some *Sideritis* species of Turkey. *Chem Nat Comp*. 39: 453-6.
- Kılıç et al., 2021 – Kılıç, Ö., Demirpolat, A., Bağcı, E., Yıldırım, Ş. (2021). Essential oil composition of *Sideritis montana* subsp. *montana*. Igdir international applied sciences congress. April 14-15, 2021. Igdir, Turkey.
- Kılıç, 2002 – Kılıç, T. (2002). *Sideritis lycia* ve *Sideritis leptoclada* türlerinin diterpen bileşiklerinin izolasyonu ve karakterizasyonu. Balıkesir Üniversitesi, Fen Bilimleri Enstitüsü, Doktora Tezi.
- Kılıç, 2003 – Kılıç, T., Yıldız, Y.K., Gören, A.C., Tümen, G., Topçu, G. (2003). Phytochemical analysis of some *Sideritis* species of Turkey. *Chem Nat Compd*. 39(5): 453-456.
- Kılıç, 2006 – Kılıç, T. (2006). Isolation and biological activity of new and known diterpenoids from *Sideritis stricta* Boiss. & Heldr. *Molecules*. 11: 257-262.
- Kılıç, 2014 – Kılıç, O. (2014). Essential Oil Composition of Two *Sideritis* L. Taxa from Turkey: A Chemotaxonomic Approach. *Asian J Chem*. 26(8): 2466-2470.
- Kılıç, 2019 – Kılıç, M. (2019). Artuklu (Mardin) Yöresinde Yetişen Bitkiler Üzerine Etnobotanik Bir Araştırma. Celal Bayar Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Botanik Bilim Dalı, Doktora Tezi.
- Kirimer et al., 2008 – Kirimer, N., Demirci, B., Işcan, G., Başer, K.H.C., Duman, H. (2008). Composition of the essential oils of two *Sideritis* species from Turkey and antimicrobial activity. *Chem Nat Compd*. 44(1): 121-123.
- Kırmızıbekmez et al., 2012 – Kırmızıbekmez, H., Arıburnu, E., Masullo, M., Festa, M., Capasso, A., Yeşilada, E. (2012). Iridoid, phenylethanoid and flavonoid glycosides from *Sideritis trojana*. *Fitoterapia*. 83: 130-6.
- Kırmızıbekmez et al., 2017 – Kırmızıbekmez, H., Karaca, N., Demirci, B., Demirci, F. (2017). Characterization of *Sideritis trojana* Bornm. Essential oil and its antimicrobial activity. *Marmara Pharm J*. 21: 860-5.
- Kirimer et al., 1992 – Kirimer, N., Koca, F., Baser, K.H.C., Ozek, T., Tanrıverdi, H., Kaya, A. (1992). Composition Of The Essential Oils Of Two Subspecies Of *Sideritis Gemanicopolitana* Bornm. *J. Essent. Oil Res*. 4: 533-534.
- Kirimer et al., 1996 – Kirimer, N., Kiirksiöbu, M., Ozek, T., Başer, K.H.C. (1996). Composition Of The Essential Oil Of *Sideritis Condensata* Boiss. Et Heldr. *Flavour And Fragrance Journal*. 1: 315-320.
- Kirimer et al., 1999 – Kirimer, N., Tabanca, N., Ozek, B., Basher, K.H.C., Tumen, G. (1999). Composition Of Essential Oils From Two Endemic *Sideritis* Species Of Turkey. *Chemistry Of Natural Compounds*. 135(1): 61-64.
- Kirimer et al., 2000 – Kirimer, N., Tabanca, N., Özek, T., Tümen, G., Baser, K.H.C. (2000). Essential Oils Of Annual *Sideritis* Species Growing In Turkey. *Pharmaceutical Biology*. 38(2): 106-111.
- Kirimer et al., 2003 – Kirimer, N., Tabanca, N., Özerk, T., Baser K.H.C. (2003). Composition of essential oils from five endemic *Sideritis* species. *J. Essent. Oil Res*. 15: 221-225.
- Kirimer et al., 2004 – Kirimer, N., Baser, K.H.C., Demirci, B., Duman, H. (2004). Essential Oils Of *Sideritis* Species Of Turkey Belonging To The Section *Empedoclia*. *Chemistry Of Natural Compounds*. 40(1): 19-23.
- Kirimer et al., 2008 – Kirimer, N., Demirci, B., Işcan, G., Baser, K.H.C., Duman, H. (2008). Composition Of The Essential Oils Of Two *Sideritis* Species From Turkey And Antimicrobial Activity. *Chemistry Of Natural Compounds*. 44(1): 121-123
- Kocabaş, Gedik, 2016 – Kocabaş, Y.Z., Gedik, O. (2016). Kahramanmaraş İl Merkezi Semt Pazarlarında Satılan Bitkiler Hakkında Etnobotanik Araştırmalar. Igdir Univ. *J. Inst. Sci. & Tech*. 6(4): 41-50.
- Koleva et al., 2002 – Koleva, I.I., Van-Beek, T.A., Linseen, J.P., De Groot, A., Eustative, L.N. (2002). Screening of extracts for antioxidant activity: a comparative study on three testing methods. *Phytochem. Anal*. 13(1): 8-17.
- Köse et al., 2010 – Köse, E.O., Deniz, İ.G., Sarıkürkçü, C., Aktaş, Ö., Yavuz, M. (2010). Chemical composition, antimicrobial and antioxidant activities of the essential oils of *Sideritis erythrantha* Boiss. and Heldr. (var. *erythrantha* and var. *cedretorum* P.H. davis) endemic in Turkey. *Food and Chem Toxicol*. 48: 2960-2965.

- Küçük et al., 2021 – Küçük, S., Kayalar, E., Kürkçüoğlu, M., Eröz-Poyraz, İ. (2021). Essential oils compositions and local names of some medicinal and aromatic plants from Lamiaceae family sold in local bazaars of Edremit-Akçay (Balıkesir-Turkey). *Biological Diversity and Conservation*. 14(3): 372-379
- Kültür, 2007 – Kültür, Ş. (2007). Medicinal plants used in Kırklareli Province (Turkey). *Journal of Ethnopharmacology*. 111: 341-364.
- Küpeli et al., 2007 – Küpeli, E., Sahin, P., Calis, I., Yeşilada, E., Ezer, N. (2007). Phenolic compounds of *Sideritis ozturkii* and their in vivo anti-inflammatory and antinociceptive activities. *J Ethnopharmacol*. 112: 356-60.
- Küpeli et al., 2007 – Küpeli, E., Şahin, F.P., Yeşilada, E., Çalış, İ., Ezer, N. (2007). In vivo antiinflammatory and antinociceptive activity evaluation of phenolic compounds from *Sideritis stricta*. *Z Naturforsch*. 62c: 519-525.
- Lee et al., 2011 – Lee, C.J., Chen, L.G., Chang, T.L., Ke, W.M., Lo, Y.F., Wang, C.C. (2011). The correlation between skin-care effects and phytochemical contents in Lamiaceae plants. *Food Chem*. 124: 833-841.
- Loğoğlu et al., 2006 – Loğoğlu, E., Arslan, S., Öktemer, A., Şakıyan, İ. (2006). Biological activities of some natural compounds from *Sideritis sipylea* Boiss. *Phytother Res*. 20: 294-297
- Loizzo et al., 2007 – Loizzo, M.R., Tundis, R., Menichini, F., Saab, A.M., Statti, G.A. (2007). Menichini F. Cytotoxic activity of essential oils from Labiatae and Lauracea families against in vitro human tumor models. *Anticancer research*. 27: 3293-3300.
- Loizzo et al., 2008 – Loizzo, M., Saab, A., Tundis, R., Menichini, F., Bonesi, M., Piccolo, V. De Cindio, B., Houghton, P. (2008). In vitro inhibitory activities of plants used in Lebanon traditional medicine against angiotensin converting enzyme (ACE) and digestive enzymes related to diabetes. *J Ethnopharmacol*. 119: 109-16.
- Menghini et al., 2005 – Menghini, L., Massarelli, P., Bruni, G., Menghini, A. (2005). Preliminary Evaluation On Anti-Inflammatory And Analgesic Effects Of *Sideritis syriaca* L. Herba Extracts. *Journal Of Medicinal Food*. 8: 227-231.
- Oymak, 2018 – Oymak, E. (2018). Bozova (Şanlıurfa) Halkının Kullandığı Doğal Bitkilerin Etnobotanik Özellikleri. Harran Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.
- Ozek et al., 1993 – Ozek, T., Baser, K.H.C., Tumen, G. (1993). The Essential Oil of *Sideritis athoa* Papanikolaou et Kokkini. *J. Essent. Oil Res*. 5: 669-670.
- Özaydın et al., 2005 – Özaydın, S., Değirmenci, T., Tümen, G., Başer, K.H.C. (2005). Plants used as analgesic in the folk medicine of Turkey. Ertuğ ZF, Proceedings of the IVth International Congress of Ethnobotany (ICEB 2005), 21-26 August 2005: İstanbul. Pp. 167-171.
- Özcan et al., 2001 – Özcan, M., Chalchat, J.C., Akgül, A. (2001). Essential oil composition of Turkish mountain tea (*Sideritis* spp.). *Food Chemistry*. 75: 459-463.
- Özdemir, Alpınar, 2015 – Özdemir, E., Alpınar, K. (2015). An ethnobotanical survey of medicinal plants in western part of central Taurus Mountains: Aladağlar (Nigde – Turkey). *Journal of Ethnopharmacology*. 166: 53-65.
- Özer, Türkmen, 2019 – Özer, H., Türkmen, N. (2019). Investigation of plants with ethnobotanical use in Gaziantep province (Turkey). *GSC Biological and Pharmaceutical Sciences*. 07(02): 071-078.
- Özer-Sağır, 2016 – Özer-Sağır, Z. (2016). Türkiye’ De Yetişen Endemik *Sideritis* L. Türlerinin (*Sideritis pisticca* Boiss. Et Heldr. Apud Benth, S. *Phrygia Bornm.*, S. *Brevibracteata* P.H. Davis) Fitokimyasal Analizleri. Balıkesir Üniversitesi, Fen Bilimleri Enstitüsü, Kimya Anabilim Dalı, Doktora Tezi.
- Özkan et al., 2001 – Özkan, M., Chalchat, J.C., Akgül, A. (2001). Essential oil composition of Turkish mountain tea (*Sideritis* spp.). *Food Chem*. 75: 459-63.
- Özkan et al., 2005 – Özkan, G., Sağdıç, O., Özkan, M., Özçelik, H., Ünver, A. (2005). Antioxidant and antibacterial activities of Turkish endemic *Sideritis* extracts. *Grasas Aceites*. 56: 16-20.
- Öztürk et al., 1996 – Öztürk, Y., Aydın, S., Öztürk, N., Başer, K.H.C. (1996). Effects of extracts from certain *Sideritis* species on swimming performance in mice. *Phytother Res*. 10: 70-73.
- Pavle et al., 2023 – Pavle, Z., Stagiopoulou, M.R., Miletić, N., Krigas, N., Lazari, D. (2023). Ecological Preferences and Diversity of Essential Oil Composition in Endangered Wild-Growing

Populations of *Sideritis sipylea* Boiss. (Lamiaceae) of the East Aegean Islands: Evidencing Antioxidant Potential, Antimicrobial and Cytotoxic Activities. *Plants*. 12: 836.

Polat et al., 2011 – Polat, R., Satıl, F., Çakılcıoğlu, U. (2011). Medicinal plants and their use properties of sold in herbal market in Bingöl (Turkey) district. *Biological Diversity and Conservation*. 4(3): 25-35.

Polat, Satıl, 2012 – Polat, R., Satıl, F. (2012). An ethnobotanical survey of medicinal plants in Edremit Gulf (Balıkesir – Turkey). *Journal of Ethnopharmacology*. 139: 626-641.

Radojevic et al., 2011 – Radojevic, I.D., Stankovic, M.S., Stefanovic, O.D., Topuzovic, M.D., Comic, L.R., Ostojic, A.M. (2011). Anti-aspergillus properties of different extracts from selected plants. *African J Mic Res*. 5: 3986-3990.

Saçlı, Akalın, 2001 – Saçlı, S., Akalın, E. (2001). Preliminary Ethnobotanical Study From Kaz Dağı (Balıkesir/Canakkale) I: Uses And Vernacular Names. *J. Fac. Pharnz. Istatzbul*. 34(2).

Sağdıç et al., 2008 – Sağdıç, O., Aksoy, A., Özkan, G., Ekici, L., Albayrak, S. (2008). Biological activities of the extracts of two endemic *Sideritis* species in Turkey. *Innovative Food Science and Emerging Technologies*. 9: 80-84.

Sara., Uğur, 2007 – Saraç, N., Uğur, A. (2007). Antimicrobial activities and usage in folkloric medicine of some Lamiaceae species growing in Muğla, Turkey. *Eurasia J Biosci*. 4: 28-37.

Sargin, 2015 – Sargin, S.A. (2015). Ethnobotanical survey of medicinal plants in Bozyazı district of Mersin, Turkey. *Journal of Ethnopharmacology*. 173: 105-126.

Sarı et al., 2010 – Sarı, A.O., Oğuz, B., Bilgiç, A., Tort, N., Güvensen, A., Şenol, S.G. (2010). Ege ve güney marmara bölgelerinde halk ilacı olarak kullanılan bitkiler. *Anadolu Ege Tarımsal Araştırma Enstitüsü Dergisi*. 20(2):1-21.

Sattar et al., 1995 – Sattar, A., Bankova, V., Kujumgiev, A., Galabov, A., Ignatova, A., Todorova, C. (1995). Chemical composition and biological activity of leaf exudates from some Lamiaceae plants. *Pharmazie*. 50(1): 62-5.

Solomou et al., 2019 – Solomou, A.D., Skoufogianni, E., Mylonas, C., Germani, R., Danalatos, N.G. (2019). Cultivation and utilization of "Greek mountain tea" (*Sideritis* spp.): current knowledge and future challenges. *Asian J Agric & Biol*. 7(2): 289-299.

Şahin, Kılıç, 2022 – Şahin, T., Kılıç, Ö. (2022). A Review: Some Plants Used for Hemorrhoids in Turkey Traditional Medicine. *Russian Journal of Biological Research*. 9(1): 8-29.

Şahin, Kılıç, 2022 – Şahin, T., Kılıç, Ö. (2022). A Review: Some Plants Used for Stomach Ailments in Turkey Traditional Medicine. *Central European Journal of Botany*. 8(1): 20-46.

Şahin, Kılıç, 2022 – Şahin, T., Kılıç, Ö. (2022). Some Plants Used for Asthma and Bronchitis in Turkey Traditional Medicine (Review). *Russian Journal of Biological Research*. 9(1): 30-44.

Şahin-Fidan, 2018 – Şahin-Fidan, E. (2018). Tek Tek Dağları Eteklerindeki Bazı Köylerde Etnobotanik Çalışma. Tezi. Harran Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans.

Tabanca et al., 2001 – Tabanca, N., Kırmır, N., Başer, K.H.C. (2001). The composition of essential oils from two varieties of *Sideritis erythrantha* var. *erythrantha* and var. *cedretorum*. *Turkish Journal of Chemistry*. 25(2): 201-208.

Tanaydın, 2021 – Tanaydın, G. (2021). Bigadiç İlçesinin (Balıkesir) Etnobotanik Özellikleri. Balıkesir Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.

Tetik et al., 2013 – Tetik, F., Civelek, S., Çakılcıoğlu, U. (2013). Traditional uses of some medicinal plants in Malatya (Turkey). *Journal of Ethnopharmacology*. 46: 331-346.

Tetik, 2011 – Tetik, F. (2011). Malatya İlinin Etnobotanik Değeri Olan Bitkileri Üzerine Bir Araştırma. Fırat Üniversitesi, Fen Bilimleri Enstitüsü, Botanik Anabilim Dalı, Yüksek Lisans Tezi.

Tevent, 2020 – Tevent, A. (2020). Çeltikçi (Burdur) İlçesinde Doğal Olarak Yayılış Gösteren Bazı Bitki Taksonlarının Etnobotanik Özellikleri. Mehmet Akif Ersoy Üniversitesi, Fen Bilimleri Enstitüsü, Biyoloji Anabilim Dalı, Yüksek Lisans Tezi.

Tunaliier et al., 2004 – Tunaliier, Z., Kosar, M., Ozturk, N., Baser, K.H.C., Duman, H., Kirimer, N. (2004). Antioxidant properties and phenolic composition of *Sideritis* species. *Chem Nat Compd*. 40: 206-10.

Turkmenoglu et al., 2015 – Turkmenoglu, F., Baysal, I., Ciftci-Yabanoglu, S., Yelekci, K., Temel, H., Pasa, S. (2015). Flavonoids from *Sideritis* species: human monoamine oxidase (hMAO) inhibitory activities, molecular docking studies and crystal structure of Xanthomicrol. *Molecules*. 20: 7454-73.

Uğur et al., 2005 – Uğur, A., Varol, O., Ceylan, O. (2005). Antibacterial activity of *Sideritis curvidens* and *Sideritis lanata* from Turkey. *Pharmaceutical Biology*. 43:47-52.

Varlıbaş-Odunkıran, 2020 – Varlıbaş-Odunkıran, Z. (2020). Hatay İlinde Etnobotanik Bir Çalışma. Yeditepe Üniversitesi, Sağlık Bilimleri Enstitüsü, Fitoterapi Ana Bilim Dalı, Yüksek Lisans Tezi.

Venditti et al., 2016 – Venditti, A., Bianco, A., Frezza, C., Serafini, M., Giacomello, G., Giuliani, C., Bramucci, M., Quassinti, L., Lupidi, G., Lucarini, D., Papa, F., Maggi, F. (2016). Secondary metabolites, glandular trichomes and biological activity of *Sideritis montana* L. subsp. *montana* from central Italy. DOI: 10.1002/cbdv.201600082.

Yeşil, İnal, 2020 – Yeşil, Y., İnal, İ. (2020). Ethnomedicinal Plants of Hasankeyf (Batman-Turkey). *Front. Pharmacol.* 11: 624710. DOI: 10.3389/fphar.2020.624710

Yeşilada et al., 1993 – Yeşilada, E., Honda, G., Sezik, E., Tabata, M., Goto, K., Ikeshiro, Y. (1993). Traditional medicine in Turkey IV. Folk medicine in the Mediterranean subdivision. *J Ethnopharmacol.* 39: 31-38.

Yeşilada et al., 1995 – Yeşilada, E., Honda, G., Sezik, E., Tabata, M., Fujita, T., Tanaka, T. (1995). Traditional medicine in Turkey. V. Folk medicine in the inner Taurus Mountains. *J Ethnopharmacol.* 46: 133-152.

Yeşilada, 2023 – Yeşilada, E. (2023). Scientific Evaluation of the Remedies Used in Turkish Folk Medicine to Treat Possible Viral Infection. *Current Traditional Medicine*. 9(6): 1-15.

Yeşilada, Ezer, 1989 – Yeşilada, E., Ezer, N. (1989). The antiinflammatory activity of some *Sideritis* species growing in Turkey. *Int J Crude Drug Res.* 27(1): 38-40.

Yücel et al., 2011 – Yücel, E., Tapırdamaz, A., Yücel-Şengün, İ., Yılmaz, G., Ak, A. (2011). Determining the usage ways and nutrient contents of some wild plants around Kisecek Town (Karaman/Turkey). *Biological Diversity and Conservation*. 4(3): 71-82.

Copyright © 2023 by Cherkas Global University



Published in the USA
Russian Journal of Biological Research
Issued from 2014
E-ISSN: 2413-7413
2023. 10(1): 30-39

DOI: 10.13187/ejbr.2023.1.30
<https://rjbr.cherkasgu.press>



Ethnobotanical Features and Pharmacological Studies of Some *Nepeta* L. Taxa

Talip Şahin ^{a, *}, Ömer Kılıç ^b

^a Department of Biology, Faculty of Arts and Sciences, Adiyaman University, Turkey

^b Department of Pharmaceutical Botanic, Faculty of Pharmacy, Adiyaman University, Turkey

Abstract

Ethnobotany is a multidisciplinary science that studies plant-human relationship and tries to understand human interaction with plants. Ethnobotanical studies aim to record the information about plants that people have passed down from generation to generation for centuries. In recent years, developments in technology, migration from village to city, modern life, relative loss of traditions and customs cause the loss of orally transmitted information about plants. In this respect, ethnobotanical studies are valuable studies that will record “humanity's knowledge of plants”.

The local population has long utilized from *Nepeta* taxa for different purposes. Most of *Nepeta* taxa have important pharmacological and biological activities and *Nepeta* taxa have many traditional usage. The majority of *Nepeta* taxa exhibit significant pharmacological characteristics and potential for pharmacy, cosmetic industry, medicine and etc.

In this study, it is aimed to review ethnobotanical and pharmacological properties of some *Nepeta* taxa to contribute to studies on the subject.

Keywords: *Nepeta*, ethnobotany, pharmacological effect, medicinal plant, aromatic plant.

1. Introduction

Since ethnobotany is a branch of science that examines the human-plant relationship, including the cultural context, it is studied by researchers from many disciplines such as botanists, pharmacists, nutritionists, agriculturalists, environmental scientists, anthropologists, political scientists, historians, geographers, economists and linguists (Ghorbani et al., 2006).

This situation allows each discipline to contribute to ethnobotany in line with their own needs. For example, pharmacists focus more on medicinal plants used by the public, while economists contribute to the registration of ethnobotanical heritage by examining plants with economic value. Today, it is possible to talk about sub-branches of ethnobotany such as archaeoethnobotany and economic botany. In addition, ethnobotany is in the intersection cluster with other disciplines researching folklore such as ethnopharmacology, ethnoecology, ethnocosmetic (Saroya, 2017).

Ethnobotanical studies have important contributions to nature, people, economy and protection of cultural heritage. In addition, it allows the identification of species that are collected intensively from nature, the identification and protection of endangered species. Recording ethnobotanical information is also very important for the discovery of new drug molecules. As in the past, plants are undoubtedly important raw material sources for the treatment of diseases today. The production of plants and herbal products is an area that affects nations economically and

* Corresponding author

E-mail addresses: talipsahin34@gmail.com (T. Şahin), omerkilic77@gmail.com (Ö. Kılıç)

socially, and also directs international relations. It enables the transfer of traditional knowledge to scientific research and thus the development of new treatment agents. It is known that many drugs used in treatment today are derived from ethnobotanical studies (Pandey and Tripathi, 2017).

It can be said that Turkey is an important center in terms of ethnobotany, thanks to its phytogeographic location and the deep-rooted cultural heritage of Anatolia. When the ethnobotanical studies conducted in Turkey are examined, it is seen that the studies mostly focus on medical uses. Although the most focused subject in the studies is the use of medicinal and food purposes, plants are frequently used for various purposes in our country such as shelter, heating, cleaning, paint, musical instruments, agricultural tools, basket-wicker making, and belief.

In this review, ethnobotanical and bioactive properties some *Nepeta* taxa are compiled, to contribute to the studies on the subject.

Table 1. Ethnobotanical and Bioactivital Properties of Some *Nepeta* taxa

Botanical name	Location of the Plant	Parts used in ethnobotany	Method of preparation in ethnobotany	Ethnobotanical uses	Reported bioactivities
<i>N. cataria</i>	Turkey (East Anatolia, Mardin), Chinese	Herb, Aerial part,	Decoction	Stomachic, stimulant, Food, Cold, Sore throat and shortness of breath, Antidepressant, Sedative, Analgesic, Bronchitis (Zhang et al., 2015; Formisano et al., 2011; Sharma et al., 2019; Altundag, Ozturk, 2011; Eksik, 2020; Özer, 2021; Kılıç, 2019; Şahin, Kılıç, 2022)	Antibacterial anti-inflammatory Aphrodisiac antimicrobial, antioxidant, bacteriostatic and fungistatic, puberty retarder (Süntar et al., 2018; Abdalla et al., 1999; Dapkevicius et al., 1998; Bourrel et al., 1993; Bernardi, 1998)
<i>N. ciliaris</i>	Uttarakhand, India	Leaf	Decoction	Flu, cough, respiratory disorders, cold and antipyretic (Joshi et al., 2016; Vohora, 1986)	Anti-inflammatory Analgesic (Hussain et al., 2012; Süntar ve ark., 2018).
<i>N. congesta</i> var. <i>cryptantha</i>	Turkey (Şanlıurfa)	Aerial part	Infusion	Headache, Cold (Abak, 2018)	-----
<i>N. crispa</i>	Iranian (Hamadan)	Aerial part	Infusion	Carminative, relaxant, sedative, restorative tonic for respiratory and nervous disorders (Sonboli et al., 2004)	Antibacterial Anti-angiogenic Antinociceptive and anti-inflammatory Anti-proliferative (Sonboli et al., 2004; Haseli et al., 2023; Badrhadad et al., 2015; Taskina et al., 2012; Zahirnia et al., 2019)
<i>N. depauperate</i>	Iranian	Whole plant	Unspecified	Wound healing and wasp, scorpion and snake bites, rheumatism (Mousavi et al., 2016)	-----
<i>N. discolor</i>	Uttarakhand - India Leh-Ladakh-	Leaf	Decoction	Leaf decoction: cold, cough and antipyretic. Decoction with honey: tuberculosis	-----

	India			(Joshi et al., 2016)	
<i>N. elliptica</i>	Uttarakhand Cemmu, Keşmir, India	Seed, Shoot, Leaf	Infusion	Digestive disorders, Food (Joshi et al., 2016) (Bhattacharjee, 2005)	Antifungal (Kumar et al., 2014)
<i>N. eriostachys</i>	Uttarakhand, Himachal Pradesh, India	Leaf, Whole plant	Unspecified	Antipyretic, Eye ailments (Joshi et al., 2016)	—————
<i>N. flavida</i>	Turkey	Aerial part	Unspecified	Flu, bronchitis and colds (Demirci, Ozhatay, 2012)	Antioxidant Herbicidal (Tepe et al., 2007; Bozok, 2019; Shakeri et al., 2016)
<i>N. floccosa</i>	Ladakh, India	Leaf	Decoction	Treat cough, cold and fever (Joshi et al., 2016)	Antioxidant (Ali et al., 2015)
<i>N. glomerulosa</i>	India	Whole plant	Unspecified	Treat itching, digestive problems and pneumonia (Nadkarni, 1976)	Antibacterial Hypnotic Effect Morphine Withdrawal Syndrome (Nezhadali et al., 2013) (Hosseini et al., 2016) (Hosseinzadeh, Ziaee, 2006)
<i>N. glutinosa</i>	Ladakh, India	Leaf	Decoction	Treat fever, pneumonia, and diarrhea (Joshi et al., 2016)	—————
<i>N. govaniana</i>	Himachal Pradesh, India	Whole plant	Decoction	Treat menstrual cramps, flu, insomnia, cold and diarrhea, Sore throat (Joshi et al., 2016)	Cytotoxicity (Dar et al., 2014)
<i>N. hindostana</i>	India	Whole plant	Decoction	Relieve sore throat, ear and toothach and fever (Nadkarni, 1976)	Effective against myocardial necrosis (provides endothelial mediator relaxation), sedative, tonic, dissolving, hepatotonic, antipyretic, lowers blood cholesterol level. Antiinflammatory Antidiabetic Antimicrobial Antioxidant Antiplatelet Dyslipidemia (Ashraf et al 1999) (Kumar et al 1998; Ahmad et al 1981; Joshi et al., 2021; Pandey et al., 2015; Siddique et al., 2018; Ansari et al., 2016; Devi, Singh, 2016; Devi, Singh, 2018)
<i>N. italica</i>	Turkey (Göller Bölgesi- Yenişarbademli Yöresi/Konya, Pertek/Tunceli, Manisa,	Aerial part, Leaf	Infusion	In addition to being used as a spice, it is used against colds, stomach flu and hair loss. As spice, As tea , Tummy ache, Common cold (Demirci, Ozhatay, 2012; Doğan, Tuzlacı, 2015; Özaydın et al.,	Analgesic Antioxidant Antimicrobial (Aydin et al 1999; Emre et al., 2011; Hasimi et al., 2015)

	Çermik/Diya rbakır)			2005; Doğanoglu et al., 2006; Başer et al., 2006; Özdemir, Alpınar, 2015)	
<i>N. laevigata</i>	Pakistan, Keşmir Ladakh, India (Uttarakhan d)	Seed, whole plant	Infusion	Dysentery, fever, colds and headaches, sore throat (Bhattacharjee, 2005)	Antioxidant Antimicrobial (Joshi, 2014; Shinwari et al., 2013)
<i>N. lagopsis</i>	Pakistan	Whole plant	Incineration (ash) and/or mist	Yara iyileştirici (Rehman et al., 2015)	_____
<i>N. leucophylla</i>	Baglund district, Nepal	Root juice, Leaf	Root juice, Paste	Cure fever (Root juice), Malaria (Paste) (Joshi et al., 2016)	Antifungal Antioxidant (Saxena, Mathela, 1996; Singh, Dhaliwal, 2018; Sharma et al., 2018; Sharma, Cannoo, 2016)
<i>N. longibractea ta</i>	Ladakh, India	Whole plant	Unspecified	Stomach ailments (Joshi et al., 2016)	Antimicrobial (Avasthi et al., 2017)
<i>N. menthoides</i>	Iranian	Unspecified	Unspecified	Febrifuge and sedative, antidepressant and anxiolytic, Carminative (Amin, 1991)	Antimicrobial Anticholinesterase Cytotoxic Anti-Inflammatory Antinociceptive Neuroprotective Memory enhancing Antioxidant (Kahkeshani et al., 2017; Ghandchi, Jamzad, 2015; Memariani et al., 2019; Sonboli et al., 2009; Süntar et al., 2018)
<i>N. nuda</i> subsp. <i>albiflora</i>	Turkey (Pertek/Tun celi)	Leaf	Unspecified	As tea (Doğan, Tuzlacı, 2015)	Antioxidant Antimicrobial (Alim et al., 2009; Mancini et al., 2009; Teber, Bursal, 2020)
<i>N. nuda</i> subsp. <i>lydia</i>	Turkey (Bingöl)	Aerial parts, Leaves	Infusion	Common cold, used as tea (Kılıç, 2016)	_____
<i>N. nuda</i> subsp. <i>glandulifera</i>	Turkey (Çivril/ Denizli)	Unspecified	Unspecified	Tummy ache (Özaydın et al., 2005)	Antioxidant Antidiabetic (Sarıkurkcu et al., 2018) (Al-Kahraman et al., 2012)
<i>N. racemosa</i>	Turkey (Erzurum)	Aerial parts	Infusion	As tea (Aksakal, Kaya, 2018)	Antimicrobial Antioxidant Anti-plasmodial (Lungoci et al., 2023; Afshar et al., 2021; Zazharskyi et al., 2020)
<i>N. septemcrena ta</i>	Egypt, Israel, Iraq, Syria and Jordan	Unspecified	Unspecified	Cardiotonic, sedative and sore throat (Tachholm, 1974)	Antimicrobial Toxicity Antipyretic Antiinflammatory Analgesic (El-Moaty, 2010; El- Hamouly, El-Hela, 2004; Moustafa et al., 2015)

<i>N. trachonitica</i>	Turkey (Pertek/Tunceli)	Aerial part	Unspecified	As tea (Doğan, Tuzlacı, 2015)	Antimicrobial Antioxidant (Köksal et al., 2017; Gökbulut, Yılmaz, 2020; Köksal et al., 2017)
<i>N. bracteata</i>	Iranian	Unspecified	Unspecified	Febrifuge and sedative, antidepressant and anxiolytic, Carminative (Amin, 1991)	Antibacterial Anti-proliferative Antiinflammatory (Süntar et al., 2018; Zhang et al., 2021)

2. Results and discussion

Continually, plant taxa have long been regarded as possessing the principal ingredients used in widely disseminated ethnomedical and pharmacological practices. Many studies have revealed that the traditional medicinal and aromatic plant taxa utilized by the local people to treat various illnesses, have not been sufficiently examined for their therapeutic/preventive potential, pharmacological, ethnobotanical and phytochemical aspects. Traditional medicinal plant taxa provide a renewable and natural source of secondary metabolites that can be exploited to create novel medications and cure a variety of human illnesses.

Many members of the Lamiaceae (Labiatae) family are widely planted for their economical qualities. *Nepeta* is among the most known genus of Lamiaceae family, include more than 285 taxa that are significant to humans as culinary, aesthetic, and medicinal plants; many *Nepeta* taxa are widely used in traditional medicine and in ethnobotany. Among many aromatic and medicinal plants, taxa belonging to the genus *Nepeta* L. are widely used in traditional medicine and have different important effects in respect to human health. Therapeutic impacts of the *Nepeta* taxa are also due to the high contents of secondary metabolites; especially flavonoids and essentials. According to the *Nepeta* taxa essential oil composition, there exist two main essential oil chemotypes (nepetolactone and 1,8-cineole and/or linalool).

Many *Nepeta* taxa have pharmacological and biological effects as sedative, antimicrobial, cytotoxic activities and anticancerous activities, apoptotic induction antioxidant, anti-inflammatory and antinociceptive effects, antidepressive, cholesterol lowering, vermifuge, antifungal, antiasthmatic, carminative, antidiabetes, acetylcholinesterase inhibition, analgesic, diuretic, diaphoretic, febrifuge, neuropharmacological activity, herbicidal, insecticidal, cardioprotective effects, angiotensin-converting enzyme inhibitory activity and insect repellent.

3. Conclusion

Due to the important effects of *Nepeta* taxa on improving the complications of various diseases, ethnobotanical and pharmacological effects the need for doing more comprehensive clinical trials for the use of *Nepeta* taxa.

Rereferences

Abak, 2018 – Abak, F. (2018). Şanlıurfa İli Lamiaceae (Ballıbabagiller) Familyasının Florası Bazı Taksonların Fitokimyasal ve Etnobotanik Özellikleri [Flora of Lamiaceae (Ballıbabagiller) Family Phytochemical and Ethnobotanical Characteristics of Some Taxa in Şanlıurfa Province]. Recep Tayyip Erdoğan Üniversitesi Fen Bilimleri Enstitüsü. Biyoloji Anabilim Dalı. Doktora Tezi. [in Turkish]

Abdalla et al., 1999 – Abdalla, A.E, Tirzite, D., Tirzitis, G., Roozen, J.P. (1999). Antioxidant activity of 1,4- dihydropyridine derivatives in β -carotene-methyl linoleate, sunflower oil and emulsions. *Food Chemistry*. 66: 1890195.

Afshar et al., 2021 – Afshar, F.H., Delazar, A., Rahimpour, Y., Navaei, N.M., Asnaashari, S., Asgharian, P. (2021). Phytochemistry and Bioactivity of *Nepeta racemosa* Lam. *Research Journal of Pharmacognosy (RJPh)*. 8(4): 1-8.

Aksakal, Kaya, 2018 – Aksakal, Ö., Kaya, Y. (2018). Erzurum ve Çevresinde Halk Tarafından Gıda Amaçlı Olarak Kullanılan Bitkiler [Plants Used for food purposes by the public in Erzurum and its surroundings]. Türkiye 10. Gıda Kongresi; 21-23 Mayıs 2008, Erzurum. [in Turkish]

- Ali et al., 2012 – Ali, T., Javan, M., Semnanian, A.S.S. (2012). Antinociceptive and anti-inflammatory activities of the essential oil of *Nepeta crispa* Willd. in experimental rat models. *Natural Product Research*. 26(16): 1529-1534.
- Ali et al., 2015 – Ali, L., Ali, S., Rizvi, T.S., Khan, A.L., Hassan, Z., Al-Harrasi, A., Hussain, J. (2015). Antioxidant Flavonoids from *Nepeta floccosa* Benth. *Rec. Nat. Prod.* 9(4): 567-571.
- Ali, Jemy, 1996 – Ali, A., Jemy, L.H. (1996). Indigenious Knowledge of Plant Use. An Ethnobotany of Chitral with Particular Reference to Medicinal Plants. Agha Khan Rural, Support Programme. Pp. 9-27.
- Alim et al., 2009 – Alim, A., Goze, I., Cetin, A., Atas, A.D., Akkus-Cetinus, S., Vural, N. (2009). Chemical composition and in vitro antimicrobial and antioxidant activities of the essential oil of *Nepeta nuda* L. subsp. *Albiflora* (Boiss.) gams. *African Journal of Microbiology Research*. 3(8): 463-467
- Al-Kahraman et al., 2012 – Al-Kahraman, Y.M.S.A., Baolch, N., Kakar, A.M., Nabi, S. (2012). In-vitro antimicrobial, insecticidal, antitumor, antioxidant activities and their phytochemical estimation of methanolic extract and its fractions of *Nepeta praetervisa* leaves. *International Journal of Phytomedicine*. 4: 531-536.
- Altundag, Ozturk, 2011 – Altundag, E., Ozturk, M. (2011). Ethnomedicinal studies on the plant resources of east Anatolia, Turkey. *Procedia Social and Behavioral Sciences*. 19: 756-777.
- Amin, 1991 – Amin, G. (1991). Popular Medicinal Plants of Iran, Research Deputy of Health Ministry, Tehran. 1: 40.
- Ansari et al., 2016 – Ansari, V.A., Arif, M., Siddiqui, H.H., Dixit, R.K. (2016). Antiplatelet activity and isolation of triterpenoidal glycoside from ethanolic extract of *Nepeta hindostana*. *Orient Pharm Exp Med*. 16: 333-337.
- Ashraf et al., 1999 – Ashraf, M.Z., Khan, M.S.Y., Hameed, H.A., Hussain, M.E., Fahim, M. (1999). Endothelium modulated vasorelaxant response of a polypharmaceutical herbal drug (lipotab) and its individual constituents. *Journal of Ethnopharmacology*. 66: 97-102.
- Avasthi et al., 2017 – Avasthi, A.S., Bhatnagar, M., Ghosal, S. (2017). High Altitude Medicinal Plants Against Multidrug Resistant Microorganisms: Screening and Chemical Profiling of the Most Active *Nepeta longibracteata*. *International Journal of Pharmacology*. 13(6): 552-562.
- Aydin et al., 1999 – Aydin, S., Demir, T., Öztürk, Y., , Başer, H.K.C. (1999). Analgesic activity of *Nepeta italica* L. *Phytotherapy Research*. 13: 20-23.
- Badrhadad et al., 2015 – Badrhadad, A., Piri, K.H., Mansouri, K. (2015). Anti-proliferative effects of some fractions of *Elaeagnus angustifolia* L. flower and aerial part of *Nepeta crispa* L. on K562 leukemic cells. *Iranian Journal of Medicinal and Aromatic Plants Research*. 31(5): 881-890.
- Başer et al., 2006 – Başer, K.H.C., Tümen, G., Malyer, H., Kırimer, N. (2006). Plants used for common cold in Turkey. *Proceedings of the IVth International Congress of Ethnobotany (ICEB 2005)*. 133-137.
- Bernardi et al., 1998 – Bernardi, M.M., Fernandes, S., Zodi, A.L., Spinosa, H.S., Gorniak, S.L. (1998). Toxic effects of catnip (*Nepeta cataria*) exposure during embryogenetic period in mice. *Toxicon*. 36: 1261-1262.
- Bhattacharjee, 2005 – Bhattacharjee, S.K. (2005). Handbook of Aromatic Plants. Pointer Publications, India, p. 311.
- Bibi et al., 2016 – Bibi, T., Ahmad, M., Edwards, S.E., Tareen, N.M., Jabeen, R., Abdullah, I. (2016). Ethnomedicinal uses of plants in the treatment of paediatric geohelminth infections in Kalat district of Northern Balochistan, Pakistan. *J. Ethnopharmacol.* 183: 176-186.
- Bozok, 2019 – Bozok, F. (2019). Herbicidal Activity of *Nepeta flavida* Essential Oil. *Journal of Essential Oil Bearing Plants*. 21(6): 1687-1693.
- Dapkevicius et al., 1998 – Dapkevicius, A., Venskutonis, R., Beek, T.A.V., Linssen, J.P.H. (1998). Antioxidant activity of extracts obtained by different isolation procedures from some aromatic herbs grown in Lithuania. *J. Sci. Food Agric*. 77: 140-146.
- Dar et al., 2014 – Dar, B.A., Lone, A.M., Qurishi, M.A. (2014). Cytotoxic activity and GC-MS analysis of the constituents of essential oil of *Nepeta govaniana* (Wall.ex Benth) from Jammu and Kashmir, India. *International Journal of Herbal Medicine*. 2(2): 58-60.
- Demirci, Ozhatay, 2012 – Demirci, S., Ozhatay, N. (2012). An ethnobotanical study in Kahramanmaras (Turkey); Wild plants used for medicinal purpose in Andırın. Kahramanmaras. *Turk. J. Pharm. Sci.* 9 (1): 75-92.

- Devi, Singh, 2010 – Devi, S., Singh, R. (2010). Antidiabetic Activity Of Methanolic Extract Of Nepeta Hindostana Herb In Streptozotocin Induced Diabetes In Rats. *International Journal of Pharmacy and Pharmaceutical Sciences*. 8(7): 330-335.
- Devi, Singh, 2017 – Devi, S., Singh, R. (2017). Assessment of lipid lowering effect of Nepeta hindostana herb extract in experimentally induced dyslipidemia. *Journal of Nutrition & Intermediary Metabolism*. 9: 17e23.
- Doğan, Tuzlacı, 2015 – Dogan, A., Tuzlacı, E. (2015). Wild Edible Plants of Pertek (Tunceli-Turkey). *Marmara Pharmaceutical Journal*. 19: 126-135.
- Doğanoğlu et al., 2006 – Doğanoğlu, Ö., Gezer, A., Yücedağ, C. (2006). Göller Bölgesi-Yenişarbademli Yöresi'nin Önemli Bazı Tıbbi ve Aromatik Bitki Taksonları Üzerine Araştırmalar [Research on Some Important Medicinal and Aromatic Plant Taxa of the Lake District-Yenişarbademli Region]. *Süleyman Demirel Üniversitesi Fen Bilimleri Enstitüsü Dergisi*. (10-1): 66-73. [in Turkish]
- Eksik, 2020 – Eksik, C. (2020). Mardin İli Artuklu, Ömerli Ve Yeşilli İlçelerinin Bazı Köylerinde Etnobotanik Çalışma [Ethnobotanical Study in Some Villages of Artuklu, Ömerli and Yeşilli Districts of Mardin Province]. *Harran Üniversitesi, Fen Bilimleri Enstitüsü, Yüksek Lisans Tezi*. [in Turkish]
- El-Hamouly, El-Hela, 2004 – El-Hamouly, M.M.A., El-Hela, A.A. (2004). Phytochemical And Biological Investigation Of The Volatile Constituents Of Nepeta Septemcrenata Ehrenb., Growing In Egypt. *Bull. Pharm. Sci., Assiut University*. 27(1): 95-98.
- El-Moaty, 2010 – El-Moaty, H.I.A. (2010). Essential oil and iridoide glycosides of Nepeta septemcrenata Erenb. *Journal of Natural Products*. 3:103-111.
- Emre et al., 2011 – Emre, I., Kurşat, M., Yılmaz, Ö., Erecevit, P. (2011). Some biological compounds, radical scavenging capacities and antimicrobial activities in the seeds of Nepeta italica L. and Sideritis montana L. subsp. montana from Turkey. *Grasas y Aceites*. 62(1): 68-75.
- Formisano et al., 2011 – Formisano, C., Rigano, D., Senatore, F. (2011). Chemical constituents and biological activities of Nepeta species. *Chem. Biodivers*. 8(10): 1783-1818.
- Ghandchi, Jamzad, 2015 – Ghandchi, S., Jamzad, M. (2015). Total Flavonoids Contents and Anti Bacterial Activity of the Extracts of two Labiateae Species: Nepeta menthoides and Thymus trautvetteri. *Journal of Medicinal Plants and By-products*. 1: 77-82.
- Ghorbani et al., 2006 – Ghorbani, A., Naghibi, F., Mosadegh, M. (2006). Ethnobotany, ethnopharmacology and drug discovery. *Iranian Journal of Pharmaceutical Sciences*. 2(2): 109-118.
- Gökbulut, Yılmaz, 2020 – Gökbulut, A., Yılmaz, G. (2020). Nepeta humilis Benth: First evaluation of phenolic profile and radical scavenging potential. *Journal of Research in Pharmacy*. 24(6). DOI: <https://doi.org/10.35333/jrp.2020.249>
- Haseli et al., 2023 – Haseli, A., Pourahmad, R., Eshaghi, M.R., Rajaei, P., Akbari-Adergani, B. (2023). Investigating the effect alcoholic extract of Nepeta crispa on the antioxidant activity and microbial and sensory properties of doogh. *Journal of Food Technology and Nutrition*. 20(25): 107-120.
- Hasimi et al., 2015 – Hasimi, N., Kizil, S., Tolan, V. (2015). Essential Oil Components, Microelement Contents and Antioxidant Effects of Nepeta italica L. and Achillea filipendulina LAM. *TEOP*. 18(3): 678-686.
- Himani et al., 2021 – Himani, M.J., Kumar, R., Prakash, O., Pant, A.K., Rawat, D.S. (2021). Chemical composition and biological activities of Nepeta hindostana (Roth) Haines, Nepeta graciliflora Benth. and Nepeta cataria L. from India. *Journal of Medicinal Herbs*. 12(2): 35-46.
- Hosseini et al., 2016 – Hosseini, A., Forouzanfar, F., Rakhshandeh, H. (2016). Hypnotic Effect of Nepeta glomerulosa on Pentobarbital-Induced Sleep in Mice. *Jundishapur J Nat Pharm Prod*. (1): e25063.
- Hosseinzadeh, Ziaee, 2006 – Hosseinzadeh, H., Ziaee, T. (2006). Effect of Nepeta glomerulosa Boiss. Aerial Parts Aqueous Extract on Morphine Withdrawal Syndrome in Mice. *Iranian Journal of Pharmaceutical Sciences Winter*. 2(1): 41-46.
- Hussain et al., 2012 – Hussain, J., Rehman, N.U., Hussain, H., Al-Harrasi, A., Ali, L., Rizvi, T.S., Ahmad, M. (2012). Analgesic, anti-inflammatory, and CNS depressant activities of new constituents of Nepeta clarkei. *Fitoterapia*. 83(3):593-598.

Joshi et al., 2016 – Joshi, R.K., Satyal, P., Setzer, W.N. (2016). Himalayan aromatic medicinal plants: a review of their ethnopharmacology, volatile phytochemistry, and biological activities. *Medicines*. 3(1):6.

Joshi, 2014 – Joshi, R.K. (2014). Antioxidant activity of essential oil of *Nepeta laevigata* (D. Don) Hand.-Mazz from Himalayan region of Uttarakhand. *American Journal of Essential Oils and Natural Products*. 2(1): 15-18.

Kahkeshani et al., 2017 – Kahkeshani, N., Hadjiakhoondi, A., Navidpour, L., Akbarzadeh, T., Safavi, M., Karimpour-Razkenari, E., Khanavi, M. (2017). Chemodiversity of *Nepeta menthoides* Boiss. & Buhse. essential oil from Iran and antimicrobial, acetylcholinesterase inhibitory and cytotoxic properties of 1,8-cineole chemotype. *Natural Product Research*. 32(22). DOI: <https://doi.org/10.1080/14786419.2017.1378202>

Kılıç, 2016 – Kılıç, Ö. (2016). An ethnobotanical survey from Bingöl (Turkey). *RA Journal of Applied Research*. 2(10); 685-691.

Kılıç, 2019 – Kılıç, M. (2019). Artuklu (Mardin) Yöresinde Yetişen Bitkiler Üzerine Etnobotanik Bir Araştırma [An Ethnobotanical Research on Plants Growing in the Artuklu (Mardin) Region]. Celal Bayar Üniversitesi Fen Bilimleri Enstitüsü Biyoloji Anabilim Dalı. Doktora Tezi. [in Turkish]

Köksal et al., 2017 – Köksal, E., Tohma, H., Kılıç, Ö., Alan, Y., Aras, A., Gülçin, I., Bursal, E. (2017). Assessment of Antimicrobial and Antioxidant Activities of *Nepeta trachonitica*: Analysis of Its Phenolic Compounds Using HPLC-MS/MS. *Sci. Pharm.* 85(24): 1-14.

Köksal et al., 2017 – Köksal, E., Tohma, H., Kılıç, Ö., Alan, Y., Aras, A., Gülçin, I., Bursal, E. (2017). Assessment of antimicrobial and antioxidant activities of *Nepeta trachonitica*: analysis of its phenolic compounds using HPLC-MS/MS. *Scientia pharmaceutica*. 85(2): 24.

Kumar et al., 2014 – Kumar, V., Mathela, C.S., Tewari, G., Singh, D. (2014). Antifungal activity of *Nepeta elliptica* Royle ex Benth. oil and its major constituent (7R)-trans,trans-nepetalactone: A comparative study. *Industrial Crops and Products*. 55: 70-74.

Lungoci et al., 2023 – Lungoci, C., Rîmbu, C.M., Motrescu, I., Serbezeanu, D., Horhoge, C.E., Vlad-Bubulac, T., Ghit, C.S., Puiu, I., Neculai-Valeanu, A.S., Robu, T. (2023). Evaluation of the Antibacterial Properties of Polyvinyl Alcohol-Pullulan Scaffolds Loaded with *Nepeta racemosa* Lam. Essential Oil and Perspectives for Possible Applications. *Plants*. 12:898.

Mancini et al., 2009 – Mancini, E., Arnold, N.A., De Feo, V., Formisano, C., Rigano, D., Piozzi, F., Senatore, F. (2009). Phytotoxic effects of essential oils of *Nepeta curviflora* Boiss. and *Nepeta nuda* L. subsp. *albiflora* growing wild in Lebanon. *Journal of Plant Interactions*. 4(4): 253-259.

Memariani et al., 2019 – Memariani, Z., Rahimi, A., Farzaei, M.H., Nejad, N.Z. (2019). *Nepeta menthoides* Boiss. & Buhse, an endemic species in Iran: A review of traditional uses, phytochemistry and pharmacology. *J Herbmed Pharmacol*. 8(3): 194-204.

Mousavi et al., 2016 – Mousavi, Z., Hosaini, A.S., Asgarpanah, J., Najafizadeh, P. (2016). Antinociceptive effect of the endemic species *Nepeta depauperata* Benth. *Jundishapur J. Nat. Pharm. Prod*. 11 (2).

Moustafa et al., 2015 – Moustafa, A.E.A., El-Azeem, H.A., Omran, M.A., Nasr, S.A.M., Abdel-Nabi, I.M., Teleb, Z.A. (2015). Biochemical Effect of *Nepeta Septemcrenata* Growing in South Sinai, Egypt. *American Journal of Ethnomedicine*. 2(3): 157-168.

Nadkarni, 1976 – Nadkarni, K.M. (1976). Indian Materia Medica. 1: 845-846. [Electronic resource]. URL: <https://archive.org/details/IndianMateriaMedicaK.M.Nadkarni/page/n383/mode/2up>

Nezhadali et al., 2011 – Nezhadali, A., Masrornia, M., Bari, H., Akbarpour, M., Joharchi, M.R., Nakhaei-Moghadam, M. (2011). Essential Oil Composition and Antibacterial Activity of *Nepeta glomerulosa* Boiss from Iran. *Journal of Essential Oil Bearing Plants*. 14(2): 241-244.

Özaydın et al., 2005 – Özaydın, S., Dirmenci, T., Tümen, G., Başer, K.H.C. (2005). Plants used as analgesic in the folk medicine of Turkey. *Proceedings of the IVth International Congress of Ethnobotany (ICEB 2005)*.

Özdemir, Alpınar, 2015 – Özdemir, E., Alpınar, K. (2015). An ethnobotanical survey of medicinal plants in western part of central Taurus Mountains: Aladaglar (Nigde – Turkey). *Journal of Ethnopharmacology*. 166: 53-65.

Özer, 2021 – Özer, D.Ö. (2021). Türkiye'nin Psikoaktif Doğal Bitki Çeşitliliğinin Etnobotanik Ve Korolojik Açidan Araştırılması [Ethnobotanical and Chorological Investigation of Turkey's Psychoactive Natural Plant Diversity]. Eskişehir Teknik Üniversitesi, Botanik Anabilim Dalı, Yüksek Lisans Tezi. [in Turkish]

Pandey et al., 2015 – Pandey, A.K., Mohan, M., Singh, P., Tripathi, N.N. (2015). Chemical Composition, Antioxidant and Antimicrobial Activities of the Essential Oil of *Nepeta hindostana* (Roth) Haines from India. *Rec. Nat. Prod.* 9(2): 224-233.

Pandey, Tripathi, 2017 – Pandey, A.K., Tripathi, Y. (2017). Ethnobotany and its relevance in contemporary research. *Journal of Medicinal Plants Studies.* 5(3): 123-129.

Raj-kumar et al., 1998 – Raj-kumar, S., Pillai, K.K., Balani, D.K., Hussain, S.Z. (1998). Antiatherosclerotic effect of Lipotab forte in cholesterol-fed rabbits. *Journal of Ethnopharmacology.* 59: 125-130.

Rehman et al., 2015 – Rehman, K., Mashwani, Z.R., Khan, M.A., Ullah, Z., Chaudhary, H.J. (2015). An ethnobotanical perspective of traditional medicinal plants from the Karak, Pakistan. *J. Ethnopharmacol.* 165: 251-259.

Salehi, Yousefzadi, 2004 – Salehi, A.S.P., Yousefzadi, M. (2004). Antimicrobial Activity and Chemical Composition of the Essential Oil of *Nepeta crispa* Willd. from Iran. *Naturforsch.* 59: 653-656.

Sarikurkcü et al., 2018 – Sarikurkcü, C., Eskici, M., Karanfil, A., Tepe, B. (2018). Phenolic profile, enzyme inhibitory and antioxidant activities of two endemic *Nepeta* species: *Nepeta nuda* subsp. *glandulifera* and *N. Cadmea*. *South African Journal of Botany.* 120: 298-301. DOI: <https://doi.org/10.1016/j.sajb.2018.09.008>

Saroya, 2017 – Saroya, A.S. (2017). Ethnobotany. New Delhi: Directorate of Knowledge Management in Agriculture. Indian Council of Agricultural Research.

Saxena, Mathela, 1996 – Saxena, J., Mathela, C.S. (1996). Antifungal activity of new compounds from *Nepeta leucophylla* and *Nepeta clarkei*. *Applied and Environmental Microbiology.* 62: 702-704.

Shakeri et al., 2016 – Shakeri, A., Khakdan, F., Soheili, V., Sahebkar, A., Shaddel, R., Asili, J. (2016). Volatile composition, antimicrobial, cytotoxic and antioxidant evaluation of the essential oil from *Nepeta sintenisii* Bornm. *Industrial Crops and Products.* 84: 224-229.

Sharma et al., 2019 – Sharma, A., Nayik, G.A., Cannoo, D.S. (2019). Pharmacology and toxicology of *Nepeta cataria* (catmint) species of genus *Nepeta*: a review. *Plant and Human Health.* (3): 285-299.

Sharma, Cannoo, 2016 – Sharma, A., Cannoo, D.S. (2016). Comparative evaluation of extraction solvents/techniques for antioxidant potential and phytochemical composition from roots of *Nepeta leucophylla* and quantification of polyphenolic constituents by RP-HPLC-DAD. *Food Measure.* 10: 658-669.

Shinwari et al., 2013 – Shinwari, Z.K., Ahmad, N., Hussain, J., Ur-Rehman, N. (2013). Antimicrobial Evaluation And Proximate Profile Of *Nepeta Leavigata*, *Nepeta Kurramensis* And *Rhynchosia Reniformis*. *Pak. J. Bot.* 45(1): 253-259.

Siddique et al., 2018 – Siddique, A.A., Gupta, P., Singh, S., Gupta, M., Misra, L., Darokar, M.P., Luqman, S., Yadav, A., Bhakuni, R.S. (2018). Antioxidant, antimicrobial activities and comparative analysis of the composition of essential oils of leaf, stem, flower and aerial part of *Nepeta hindostana*. *Official Journal of the Societa Botanica Italiana.* 153(2): 242-249. DOI: <https://doi.org/10.1080/11263504.2018.1473301>

Singh, Dhaliwal, 2018 – Singh, J., Dhaliwal, A.S. (2018). Novel Green Synthesis and Characterization of the Antioxidant Activity of Silver Nanoparticles Prepared from *Nepeta leucophylla* Root Extract. *Analytical Letters.* 52(2): 213-230. DOI: <https://doi.org/10.1080/00032719.2018.1454936>

Sonboli et al., 2009 – Sonboli, A., Gholipour, A., Yousefzadi, M., Mojarad, M. (2009). Antibacterial Activity and Composition of the Essential Oil of *Nepeta menthoides* from Iran. *Natural Product Communications.* 4(2): 283-286.

Sonbolia et al., 2004 – Sonbolia, A., Salehi, P., Yousefzadi, M. (2004). Antimicrobial activity and chemical composition of the essential oil of *Nepeta crispa* Willd. from Iran. *Z. Naturforsch.* 59(9-10): 653-656.

- Süntar et al., 2018 – Süntar, I., Nabavi, S.M., Barreca, D., Fischer, N., Efferth, T. (2018). Pharmacological and chemical features of Nepeta L. genus: Its importance as a therapeutice agent. *Phytotherapy Research*. 32(2): 185-198.
- Şahin, Kılıç, 2022 – Şahin, T., Kılıç, Ö. (2022). A Review: Some Plants Used for Stomach Ailments in Turkey Traditional Medicine. *Central European Journal of Botany*. 8(1): 20-46.
- Tachholm, 1974 – Tachholm, V. (1974). Student's Flora of Egypt, second ed. Cairo University, Coop, Beirut. Talebi S.M., Nohooji M.G., Yarmohammadi M., 2020. Essential oil analysis of eight Nepeta taxa in Iran. *Mediterr. Bot.* 41(1): 43-53.
- Teber, Bursal, 2020 – Teber, İ., Bursal, E. (2020). Phenolic Compounds and Antioxidant Activity of Nepeta Nuda Subsp. Albiflora. *International Letters of Natural Sciences*. 79: 1-8.
- Tepe et al., 2007 – Tepe, B., Daferera, D., Sihoğlu-Tepe, A., Polissiou, M., Sökmen, A. (2007). Antioxidant activity of the essential oil and various extracts of Nepeta flavida Hubç-Mor. From Turkey. *Food Chemistry*. 103: 1358-1364.
- Vohora, 1986 – Vohora, S.B. (1986). Unani joshandah drugs for common cold, catarrh, cough, and associated fevers. *J. Ethnopharmacol.* 16 (2–3): 201-211.
- Zahirnia et al., 2019 – Zahirnia, A., Boroomand, M., Nasirian, H., Soleimani-Asl, S., Salehzadeh, A., Dastan, D. (2019). The cytotoxicity of malathion and essential oil of Nepeta crispa (lamiales: lamiaceae) against vertebrate and invertebrate cell lines. *Pan African Medical Journal*. 33: 285.
- Zazharskyi et al., 2020 – Zazharskyi, V.V., Davydenko, P.O., Kulishenko, O.M., Borovik, I.V., Zazharska, N.M., Brygadyrenko, V.V. (2020). Antibacterial and fungicidal activities of ethanol extracts of 38 species of plants. *Biosyst. Divers.* 28(3): 281-289.
- Zhang et al., 2015 – Zhang, L., Wang, Y., Yang, D., Zhang, C., Zhang, N., Li, M., Liu, Y. (2015). Platycodongrandiflorus. An ethnopharmacological, phytochemical and pharmacological review. *J. Ethnopharmacol.* 164: 147-161.
- Zhang et al., 2021 – Zhang, M., Chen, M., Hou, Y., Fan, C., Wei, H., Shi, L., Ma, G., Zhang, J. (2021). Inflammatory and Cytotoxic Activities of Abietane Terpenoids from Nepeta bracteata Benth. *Molecules*. 26: 5603. DOI: <https://doi.org/10.3390/molecules26185603>